



RESEARCH

Open Access

Genome-wide RNAi screen reveals the E3 SUMO-protein ligase gene *SIZ1* as a novel determinant of furfural tolerance in *Saccharomyces cerevisiae*

Han Xiao¹ and Huimin Zhao^{1,2*}

Abstract

Background: Furfural is a major growth inhibitor in lignocellulosic hydrolysates and improving furfural tolerance of microorganisms is critical for rapid and efficient fermentation of lignocellulosic biomass. In this study, we used the RNAi-Assisted Genome Evolution (RAGE) method to select for furfural resistant mutants of *Saccharomyces cerevisiae*, and identified a new determinant of furfural tolerance.

Results: By using a genome-wide RNAi (RNA-interference) screen in *S. cerevisiae* for genes involved in furfural tolerance, we identified *SIZ1*, a gene encoding an E3 SUMO-protein ligase. Disruption of *SIZ1* gene function by knockdown or deletion conferred significantly higher furfural tolerance compared to other previously reported metabolic engineering strategies in *S. cerevisiae*. This improved furfural tolerance of *siz1Δ* cells is accompanied by rapid furfural reduction to furfuryl alcohol and leads to higher ethanol productivity in the presence of furfural. In addition, the *siz1Δ* mutant also exhibited tolerance towards oxidative stress, suggesting that oxidative stress tolerance related proteins may be under the SUMO regulation of SIZ1p and responsible for furfural tolerance.

Conclusions: Using a genome-wide approach, we identified a novel determinant for furfural tolerance, providing valuable insights into the design of recombinant microbes for efficient lignocellulose fermentation.

Keywords: Furfural tolerance, RAGE, *Saccharomyces cerevisiae*, *SIZ1*, SUMO E3 ligase

Background

There is a growing interest worldwide in using lignocellulose, the most abundant renewable biomass, to replace cereal substrates in the production of biofuels and biochemicals [1,2]. However, efficient fermentation of lignocellulosic hydrolysates is limited by inhibitors that are inevitably released during pretreatment and hydrolysis of lignocellulosic substrates [3]. Formed by dehydration of pentoses during dilute acid pretreatment of lignocelluloses, furfural is one of the major inhibitors present in lignocellulosic hydrolysates [4]. The toxicity of hydrolysates correlates with furfural concentration, with 1 to 5 g/L

of furfural leading to complete growth inhibition of *Escherichia coli*, *Zymomonas mobilis* and *Saccharomyces cerevisiae*, significantly reducing the yield and productivity of desired products [5-8]. Although physical or chemical strategies for furfural detoxification can be adopted during fermentation, the additional equipment and time required increase the production costs [9]. Thus, improving furfural tolerance in microorganisms would provide a cost-effective means for lignocellulose fermentation.

S. cerevisiae is the most widely studied model organism for furfural tolerance and has higher furfural tolerance compared to other potential biofuel and biochemical production hosts [10-12]. Furfural modulates expression of genes involved in a variety of general stress responses in *S. cerevisiae*, including oxidative stress, nutrient starvation, DNA damage, unfolded protein response, as well as osmotic and salt stress [11]. However, whether and how

* Correspondence: zhao5@illinois.edu

¹Department of Chemical and Biomolecular Engineering, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

²Departments of Chemistry, Biochemistry, and Bioengineering, Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

these genes contribute to furfural tolerance is unknown [13]. To date, the known mechanism of furfural detoxification is its reduction into the less toxic furfuryl alcohol through reduced nicotinamide adenine dinucleotide phosphate (NADPH)-dependent enzymes [13,14]. Attributed to the significant increase in mRNA abundance and protein expression level observed in adapted *S. cerevisiae* under stress challenge, the NADPH-dependent oxidoreductases ADH7p and YKL071Wp were found to be responsible for furfural detoxification [15,16]. Due to limited knowledge on the mechanisms of furfural toxicity towards cells, strategies for improving furfural tolerance focus mainly on overexpression of the enzymes that convert furfural to furfuryl alcohol [12,17]. To fill this knowledge gap and at the same time develop strains with strong furfural resistance, genetic determinants of furfural tolerance need to be identified.


Genes associated with furfural tolerance have been identified by comparative analyses of wild-type strains with furfural tolerant mutants generated via random mutation, directed evolution or adaptation strategies. However, the existence of multiple simultaneous mutations in these tolerant strains often complicates the analyses [18]. Therefore, dissecting the functional contribution of each gene towards furfural tolerance remains a significant challenge. RAGE (RNAi-assisted genome evolution) is a recently developed genome engineering method that can continuously improve a desired trait by allowing the sequential introduction of tractable reduction-of-function

modifications to the genome [19]. In this study, we used RAGE to select for clones with increased furfural tolerance. The genome-wide RNAi library of *S. cerevisiae* BY4741 was selected for clones with increased furfural tolerance to discover determinants of furfural resistance. Deletion of the gene *SIZ1*, which encodes an E3 SUMO-protein ligase, was found to play an important role in tolerance to furfural and general oxidative stress in *S. cerevisiae*.

Results

RAGE screen and isolation of furfural resistant strains

To uncover new genetic determinants of furfural tolerance, we sought to determine if furfural tolerance can be enhanced through reduction of gene function by using RAGE to select for furfural resistant mutants (Figure 1A). Reconstitution of the RNAi machinery in *S. cerevisiae* BY4741 was carried out as previously reported [19] to yield the BAD strain. The genomic DNA derived RNAi library was constructed with additional modifications to prevent self-ligation of vectors and fragments [20]. DNA sequencing of 17 randomly picked plasmids from the RNAi library showed that only one locus was targeted by each RNAi construct (Additional file 1: Figure S1). The number of *Sau3AI* digested fragments (35,837) from *S. cerevisiae* genomic DNA [21] was considered as the number of possible equiprobable variants. With a library size of 3.4×10^5 , more than 99% coverage of the yeast genome was achieved [22].



By selecting mutant colonies that grew larger than that of strain BAD-P (strain BAD with plasmid backbone alone) on synthetic complete medium deficient in uracil (SC-URA) plates containing 0.8 g/L furfural, we isolated and confirmed four clones with increased furfural tolerance. Sequencing revealed that three out of the four RNAi constructs from these furfural resistant clones contained the same fragment of the *SIZ1* gene, which encodes an E3 small ubiquitin-like modifier (SUMO)-protein ligase (Additional file 1: Figure S2). The RNAi construct isolated from the fourth clone contained a fragment of the *GCN4* gene encoding a basic leucine zipper transcriptional activator of amino acid biosynthesis (Additional file 1: Figure S3). Strains expressing the RNAi cassettes targeting genes *SIZ1* (*SIZ1*-kd) and *GCN4* (*GCN4*-kd) exhibited significant improvement in furfural tolerance compared to the control BAD-P strain (Figure 1B and C). Notably, *SIZ1*-kd and *GCN4*-kd strains showed no growth advantage over the control strain in the absence of furfural (Additional file 1: Table S3).

Disruption of *SIZ1* function increased furfural tolerance in *S. cerevisiae*

To determine if the reduction or loss of *SIZ1* and *GCN4* gene function contributes to furfural resistance, furfural tolerance of the respective knockout mutants was tested. The maximum specific growth rate in the presence of 0.8 g/L furfural of the *siz1Δ* strain was 73% higher than wild type. Increased furfural tolerance was also found in the *siz1Δ* strain when higher furfural concentrations were adopted (Additional file 1: Table S4). Complementation of the *siz1Δ* strain with a plasmid-borne copy of gene *SIZ1* but not with plasmid alone (*siz1Δ*-P) restored furfural sensitivity (Table 1). On the other hand, deletion of *GCN4* did not phenocopy the improved tolerance observed for the *GCN4*-kd strain, suggesting that gene dosage is important for *GCN4* associated furfural tolerance (Table 1). We chose to focus on *SIZ1* as *SIZ1*-kd and *siz1Δ* have a more significant effect on furfural tolerance compared to *GCN4*-kd.

To investigate whether increased furfural tolerance via deletion of *SIZ1* is a strain-specific or general attribute,

SIZ1 was also deleted in two other *S. cerevisiae* strains: HZ848 [23] and W303a [24]. Furfural tolerance of these mutants was tested in the presence of 0.8 g/L furfural. Strains HZ848-*siz1Δ* and W303a-*siz1Δ* exhibited 27% and 58% higher maximum specific growth rates respectively, as compared to their respective parent strains (Table 2). These results showed that *SIZ1* was indeed an important determinant for furfural resistance in *S. cerevisiae*.

Increased rate of furfural reduction and ethanol productivity by *siz1Δ* strain

Having demonstrated that disruption of *SIZ1* gene function greatly increases furfural tolerance, we sought to determine the effect and utility of the enhanced furfural tolerance observed for the *siz1Δ* strain. Batch fermentation containing 20 g/L glucose and 0.8 g/L furfural was conducted using the *siz1Δ* and wild type (BAD) strains. While both control and *siz1Δ* strains experience a delay in entering exponential growth in the presence of furfural, a shorter initial lag was observed in the *siz1Δ* strain (Figure 2A). Finally, furfural was consumed and converted to the less toxic furfuryl alcohol at a rate that was 48% faster in the *siz1Δ* strain compared to that of the wild type (0.031 g/(L·h) versus 0.021 g/(L·h)) (Figure 2B). Strain *siz1Δ* consumed all glucose in 30 h, which was 18 h faster than that of the wild-type strain BAD (Figure 2B and C). As a result, strain *siz1Δ* was able to produce 9.0 g/L ethanol after 30 h, resulting in 275% higher productivity and 254% higher ethanol yield than that observed for strain BAD (Figure 2C, Additional file 1: Table S5). The molar ratios of carbon used for ethanol production were comparable between strain BAD and Δ *siz1*, indicating the improved furfural tolerance in strain Δ *siz1* was not at the cost of ethanol yield (Additional file 1: Table S5). Overall, these results demonstrate that the increased furfural tolerance observed with disruption of *SIZ1* function was accompanied by faster furfural reduction and this improved trait has clear utility in improving the efficiency of lignocellulose fermentation containing furfural.

Table 1 Furfural tolerance assay of *siz1Δ* mutant and its complementary derivatives

Strain	Maximum specific growth rate (h ⁻¹)
BAD	0.15 ± 0.00
<i>siz1Δ</i>	0.26 ± 0.00
<i>siz1Δ</i> -P	0.26 ± 0.01
<i>siz1Δ</i> - <i>SIZ1</i>	0.19 ± 0.00
<i>gcn4Δ</i>	0.16 ± 0.01

Strains were grown in the presence of 0.8 g/L furfural. Results are presented as mean ± SD (n = 3).

Table 2 Furfural tolerance assay of *S. cerevisiae siz1Δ* mutants in SC medium containing 0.8 g/L furfural

Strain	Maximum specific growth rate (h ⁻¹)
HZ848	0.15 ± 0.01
HZ848- <i>siz1Δ</i>	0.19 ± 0.00
W303a	0.19 ± 0.01
W303a- <i>siz1Δ</i>	0.30 ± 0.01

The maximum specific growth rates of different *S. cerevisiae siz1Δ* mutants are statistically significant over their corresponding wild-type ($P < 0.05$) as determined by the Student *t*-test. Results are presented as mean ± SD (n = 3).




Figure 2 Growth and metabolite profiles in batch fermentation of parent BAD and *siz1Δ* strains. Strains were grown in SC medium with 20 g/L glucose in the presence and absence of 0.8 g/L furfural. **(A)** Cell growth as measured by optical density (OD)₆₀₀. **(B)** Furfural consumption and furfuryl alcohol production. **(C)** Glucose consumption and ethanol production. Error bars represent SD of the mean (n = 3).

Comparison of *SIZ1* deletion with other previously reported strategies for improving furfural tolerance in *S. cerevisiae*

The rapid furfural reduction observed for *siz1Δ* cells is reminiscent of furfural detoxification by enzymes that catalyze aldehyde reduction coupled with cofactors NADPH and/or NADH [13]. Indeed, overexpression of various aldehyde reduction enzymes encoded by genes *YKL071W*, *ALD6*, *ADH7* and *ARI1* have been demonstrated to be strongly associated with furfural resistance in yeast [8,12,17]. In addition, overexpression of glucose-6-phosphate dehydrogenase encoding gene *ZWF1* and transcriptional activator encoding gene *MSN2*, which are involved in regeneration of NAD(P)H and stress response, respectively, have also been confirmed to increase furfural tolerance in *S. cerevisiae* [25,26]. To compare these reported targets with *siz1Δ*-associated furfural tolerance, individual overexpression of each gene was performed in strain BAD. Unexpectedly, only overexpression of gene *ADH7* and *ARI1* resulted in increased furfural tolerance (Figure 3), which may be attributed to the different promoters adopted for over-expression, different growth media tested for furfural tolerance and/or




Figure 3 Maximum specific growth rates to furfural tolerance assay of parent strain BAD and its indicated derivatives in SC medium containing 0.8 g/L furfural. The Student *t*-test was performed to determine whether the specific growth rates of indicated derivatives was statistically significant over that of the parent strain BAD. **P* < 0.05; ***P* < 0.01. Error bars represent the SD of the mean (n = 3).

different strains used. Among the engineered strains tested, the *siz1Δ* mutant exhibited the highest maximum specific growth rate in the presence of 0.8 g/L furfural (Figure 3).

Furfural tolerance is specific to SIZ1p and not to other SUMO E3 ligases

Protein sumoylation, an important post-translational modification in various cellular processes, involves the covalent attachment of the SUMO polypeptide to specific lysine residues of target proteins [27]. The E3 SUMO-protein ligase facilitates the transfer of SUMO to the substrate proteins [28]. To investigate whether furfural tolerance is specific to SIZ1p, genes *SIZ2*, *MMS21* and *CST9* that encode for the other three E3 SUMO-protein ligases in yeast [29] were individually deleted but did not affect furfural tolerance of the cells (Figure 4).

As downregulation of gene *GCN4* increased the furfural tolerance of strain BAD (Figure 1), this strategy was evaluated in *siz1Δ* strain. No significant difference in the maximum specific growth rates was found between strain *siz1Δ-GCN4-kd* and the *siz1Δ* strain in the presence of 0.8 g/L furfural. This observation suggests that downregulation of *GCN4* may act in the same pathway as *SIZ1* deletion (Figure 5).

Disruption of SIZ1 function increases tolerance to oxidative stresses

According to a previous study, downregulation of *GCN4* increases tolerance of a furfural-like chemical 5-hydroxymethylfurfural (5-HMF), which is another major inhibitor in lignocellulose hydrolysates that is derived from dehydration of hexoses in lignocellulosic hydrolysates [15,30]. The *siz1Δ* mutant also exhibits higher maximum specific




Figure 4 Furfural tolerance assay of indicated E3 small ubiquitin-like modifier (SUMO)-protein ligase gene deletion mutants in SC medium containing 0.8 g/L furfural. Error bars represent SD of the mean (n = 3).





Figure 5 Furfural tolerance assay of *siz1Δ* strain and its derivatives in SC medium containing 0.8 g/L furfural. Error bars represent the SD of the mean (n = 3).

growth rate in the presence of 1.26 g/L 5-HMF compared to the control strain (Figure 6A), while the maximum specific growth rates of both strains were similar in the absence of 5-HMF (Additional file 1: Table S3). Given the higher tolerance achieved by *SIZ1* deletion as compared to downregulation of *GCN4* in the presence of either furfural or 5-HMF, other proteins that are under the SUMO regulation of SIZ1p may also be involved in furfural tolerance besides *GCN4p* (Figure 1C, Table 1, Figure 6A and Figure 5). Furfural induces accumulation of reactive oxygen species (ROS), the toxicity of which is greatly attenuated in hosts with strong oxidative stress tolerance [10]. To investigate whether *siz1Δ* strain has a detoxification effect on ROS, the oxidative stress tolerance of the *siz1Δ* mutant was further tested in the presence of 1.72 mg/L menadione, which is known to generate ROS *in vivo* [31]. As shown in Figure 6B, the maximum specific growth rate of the *siz1Δ* mutant was 28% higher than that of the wild type. This result indicated that proteins that are related to oxidative stress tolerance may be under the SUMO regulation of *SIZ1* and responsible for furfural tolerance.

Discussion

Furfural toxicity is a major hurdle in the economical fermentative processes for biofuel and biochemical production using lignocellulosic hydrolysates as substrate [32]. Given the incomplete knowledge on furfural tolerance mechanisms, discovering new targets of furfural resistance would facilitate development of new metabolic engineering strategies for improving furfural tolerance. Whereas significant attention has been paid to the induced



genes in furfural tolerance studies, the importance of the repressed genes is often neglected [13]. In this study, RAGE was used for selection of furfural resistant mutants, in order to identify those genes with previously undiscovered roles in furfural tolerance. Genes with downregulation and/or loss of function can be selected out in our case, which distinguishes RAGE from other screening methods (for example, gain-of-function-based screening and *S. cerevisiae* single gene-knockout collection-based screening). In a previous study, RAGE was demonstrated to continuously improve acetic acid tolerance by accumulating reduction-of-function modifications in the genome [19]. In this study, however, no further improvement of furfural tolerance was observed after the second round of selection in the *siz1Δ* strain by RAGE (data not shown).

RNAi cassettes targeting genes *SIZ1* and *GCN4* were recovered during selection for furfural resistance and were shown to increase furfural tolerance (Figure 1). Downregulation of *GCN4* also increased 5-HMF tolerance

according to a previous study [15]. *GCN4*p is a transcriptional activator of gene expression related to amino acid biosynthesis during amino acid starvation in yeast [33,34]. Downregulated expression of *GCN4* could be an efficient means of energy utilization for economic pathway development [15]. However, deletion of *GCN4* did not show increased furfural tolerance in our work, suggesting that an appropriate expression level of amino acid biosynthetic genes may facilitate cell survival under stress challenge.

Replicated discovery of *SIZ1* by RAGE in the presence of furfural, along with the ability of *siz1Δ* to increase furfural tolerance in different *S. cerevisiae* strains, showed the important role of *SIZ1* in furfural resistance (Figure 1, Table 2, Additional file 1: Figure S2). To our knowledge, this is a novel determinant of furfural tolerance. SUMO-modified proteins participate in transcription, nuclear transport, cell cycle, DNA repair and signal transduction [35]. The vast majority of sumoylation in yeast is mediated by SUMO E3 ligase [36]. *SIZ1*p, together with another SUMO E3 ligase *SIZ2*p, accounts for 90% of the total sumoylation in yeast [37,38]. As furfural tolerance is *siz1Δ*-dependent (Figure 4), targets that improve furfural tolerance may be exclusively under the SUMO regulation of *SIZ1*p. Alternatively, these targets may also be under SUMO regulation of other E3 SUMO-protein ligases, but the effect of furfural tolerance from deletion of other E3 SUMO-protein ligases was masked by interactions with other targets which increased susceptibility to furfural.

Downregulation of *GCN4* and deletion of *SIZ1* increased furfural tolerance (Figure 1 and Figure 2). However, no increase of furfural tolerance was found when *GCN4* expression was reduced in the *siz1Δ* strain (Figure 5), indicating these two genes may act in the same pathway. *GCN4*p stability was demonstrated to be regulated by sumoylation, which occurs after it binds to target promoters and facilitates the subsequent removal of *GCN4*p from these promoters to ensure accurate transcription of its target genes [39,40]. A possible mechanism for furfural tolerance in the *siz1Δ* strain was speculated upon here. In the *siz1Δ* strain, non-sumoylated *GCN4*p cannot dissociate from target promoters after recruitment of RNA polymerase II, which may decrease the transcriptional efficiency of the target genes and facilitate cell survival in the presence of furfural.

SIZ1-kd or deletion strains exhibited higher furfural tolerance as compared to the *GCN4*-kd strain, indicating more SUMO targets of *SIZ1*p are likely to be involved in furfural resistance besides *GCN4*p (Figure 1 and Table 1). A newly identified SUMO substrate of *SIZ1*p is the NADHX dehydratase YKL151Cp, which converts (S)-NADHX to NADH [41,42]. More reducing power for furfural detoxification may be generated through this reaction. In addition, the *siz1Δ* strain also exhibited increased oxidative stress tolerance, suggesting that

determinants of oxidative stress tolerance may be protein substrates of SIZ1p responsible for furfural tolerance (Figure 6B). A total of 159 proteins were identified to be sumoylated in a proteomics study [40], among which the 6-phosphogluconate dehydrogenase GND1p, the basic leucine zipper transcription factor SKO1p and the redoxin peroxidase TSA1p are implicated in oxidative stress response in yeast [43-45]. These candidates could be possible determinants of furfural resistance under SUMO regulation of SIZ1p. Identification of protein substrates that are differentially sumoylated in wild type and *siz1Δ* cells in the presence of furfural will further reveal the detailed molecular mechanism of furfural resistance in the mutant.

Conclusions

In this study, RNAi knockdown of genes *SIZ1* and *GCN4* was demonstrated to improve furfural tolerance in *S. cerevisiae*. The *siz1Δ* mutant was further found to exhibit superior performance with cell growth, glucose consumption, furfural consumption and ethanol productivity as compared to the parent strain, while the *gcn4Δ* strain did not exhibit improved furfural tolerance. Deletion of *SIZ1* also resulted in higher furfural tolerance in different *S. cerevisiae* strains, indicating *SIZ1* deletion may play an important role in furfural resistance in *S. cerevisiae* strains. To our knowledge, this is a novel determinant of furfural resistance. Preliminary exploration of furfural tolerance in the *siz1Δ* mutant showed that the proteins responsible for furfural tolerance, among which GCN4p is a possible candidate, may be exclusively under the SUMO regulation by SIZ1p. Besides furfural tolerance, the *siz1Δ* mutant also exhibited tolerance towards oxidative stress, suggesting that proteins that are related to oxidative stress tolerance may be under the SUMO regulation of SIZ1p and responsible for furfural tolerance. These findings provide valuable insights into the engineering of furfural resistant microbes for efficient lignocellulose-based fermentation.

Methods

Strains and growth media

The strains and plasmids used in this study are listed in Table 3. Cells were grown in liquid SC [46], SC-URA or synthetic complete medium deficient in leucine (SC-LEU) supplemented with 20 g/L glucose as the carbon source or on solid 1% yeast extract, 2% peptone, 0.01% adenine hemisulfate, 2% glucose and 2% agar (YPAD) medium unless otherwise noted. The initial pH value of SC medium was adjusted to 5.6 using 12 M NaOH.

Construction of plasmids, genome-wide RNAi library and reconstitution of RNAi machinery in *S. cerevisiae* BY4741

The primers used in this study are listed in Additional file 1: Table S1. Plasmid constructions are summarized

in Additional file 1: Table S2. All plasmid construction was performed by In-fusion HD cloning (Clontech Laboratories, Inc., Mountain View, CA, USA) following the manufacturer's instructions, or by the DNA assembler method [23]. Construction of the genomic library of *S. cerevisiae* BY4741 was carried out as previously described [19] with modifications to prevent self-ligation of vectors and fragments [20]. Finally, a library size of 2.6×10^6 transformants was obtained, while the control reaction with only linearized plasmid gave 4×10^4 transformants. The plasmid library was isolated from an overnight *E. coli* culture.

DNA transformation of *S. cerevisiae* strains

DNA transformation of *S. cerevisiae* strains was carried out using the method developed by Gietz and Schiestl [49].

RAGE screen for increased furfural tolerance

The RNAi library (20 μg) or control plasmid pRS416-TTrcx was transformed into the BAD strain harboring the RNAi machinery. A library size of 3.4×10^5 was achieved, ensuring >99% coverage of the yeast genome [22]. Following transformation, yeast cells were recovered in 1 mL YPAD medium for 4 h, washed with ddH₂O and plated onto solid SC-URA medium containing 0.8 g/L furfural. The library and control plates were incubated at 30°C for 3 to 5 days. Thirty-three colonies of sizes bigger than the largest colonies on the control plates were picked from the library plates into SC-URA liquid medium. The growth performance of the selected colonies and control strain were compared in the presence of 0.8 g/L furfural. The initial OD₆₀₀ for all the strains was 0.2, and the growth rate was measured after 24 h. The RNAi plasmids from the top 14 strains with OD₆₀₀ values at least 20% higher than the control strain were isolated and amplified in *E. coli*. The selected plasmids were then individually retransformed, of which four were able to retain the enhanced furfural tolerance in a fresh genetic background with three biological replicates. The four plasmids were sequenced with the primer pRS416-TTrc-S (Additional file 1: Table S1).

Spot assay

Yeast cells in the stationary phase were transferred into 5 mL of SC media in a 15-mL round-bottom Falcon tube at an initial OD₆₀₀ of 0.2 and grown to an OD₆₀₀ of 0.7 (30°C, 250 rpm). The cells were serially diluted 10-fold with sterile water and 5 μl of each dilution was spotted onto furfural-free SC agar (control) and SC agar medium containing 0.8 g/L furfural. The plates were incubated at 30°C for 2 to 5 days.

Table 3 Strains and plasmids used in this study

Strains or plasmids	Characteristics	Reference or source
Strains		
<i>S. cerevisiae</i>		
BY4741	<i>MATa his3Δ0 leu2Δ0 met15Δ0 ura3Δ0</i>	[47]
BAD	BY4741/Δ::TEF1p-ago1-TPI1p-dcr1	This study
BAD-P	BAD/pRS416-TTrcx	This study
<i>SIZ1</i> -kd	BAD/pRS416-TTrcx-siz1	This study
<i>GCN4</i> -kd	BAD/pRS416-TTrcx-gcn4	This study
<i>siz1Δ</i>	BAD/ <i>siz1Δ</i> ::leu2	This study
<i>siz1Δ</i> -P	<i>siz1Δ</i> /pRS416e	This study
<i>siz1Δ</i> - <i>SIZ1</i>	<i>siz1Δ</i> /pRS416e-siz1	This study
<i>gcn4Δ</i>	BAD/ <i>gcn4Δ</i> ::leu2	This study
<i>siz2Δ</i>	BAD/ <i>siz2Δ</i> ::leu2	This study
<i>mms21Δ</i>	BAD/ <i>siz1Δ</i> ::leu2	This study
<i>cst9Δ</i>	BAD/ <i>cst9Δ</i> ::leu2	This study
BAD-YKL071W	BAD/pRS416e-ykl071w	This study
BAD-ZWF1	BAD/pRS416e-zwf1	This study
BAD-MSN2	BAD/pRS416e-msn2	This study
BAD-ALD6	BAD/pRS416e-ald6	This study
BAD-ADH7	BAD/pRS416e-adh7	This study
BAD-ARI1	BAD/pRS416e-ari1	This study
<i>siz1Δ</i> - <i>GCN4</i> -kd	<i>siz1Δ</i> /pRS416-TTrcx-gcn4	This study
HZ848	<i>MATa, ade2-1, Δura3, his3-11, 15, trp1-1, leu2-3, 112, and can1-100</i>	[23]
HZ848- <i>siz1Δ</i>	HZ848/ <i>siz1Δ</i> ::ura3	This study
W303a	<i>MATa; ura3-1; trp1Δ 2; leu2-3,112; his3-11,15; ade2-1; can1-100</i>	[24]
W303a- <i>siz1Δ</i>	W303a/ <i>siz1Δ</i> ::hygromycin B	This study
<i>E. coli</i>		
DH5α	General cloning host	Takara
WM1788	Cloning host	Provided by Professor William Metcalf
Plasmids		
pRS416	Yeast centromere with URA3 marker	[48]
pRS425-TEF1p-Pmel-PGK1t	Yeast gene expression vector	[19]
pRS416e	Derived from pRS416, with <i>TEF1</i> promoter and <i>PGK1</i> terminator added	This study
pRS-delta-KanMX-LoxP-TEF1p-AGO1-PGK1t-TPI1p-DCR1-GPD1t	Helper plasmid for integration of <i>S. castellii</i> RNAi pathway into delta-site	[19]
pRS416-TTrc	Derived from pRS416, with convergent promoters to produce dsRNA	[19]
pRS416-TTrcx	Derived from pRS416-TTrc, with <i>XhoI</i> restriction recognition sequence instead of <i>BamHI</i>	This study
pRS416-TTrcx-siz1	Derived from pRS416-TTrcx, with gene <i>SIZ1</i> fragment added	This study
pRS416-TTrcx-gcn4	Derived from pRS416-TTrcx, with gene <i>GCN4</i> fragment added	This study
pRS415	Yeast centromere with LEU2 marker	[48]
pUG6	The loxP-KanMX-loxP disruption module	Euroscarf
pUG72	The loxP-URA3-loxP disruption module	Euroscarf
pLHCX	Template for amplification of hygromycin B resistance gene	Clontech

Table 3 Strains and plasmids used in this study (Continued)

pXZ5	Derived from pUG72, with hygromycin B resistance gene expression cassette instead of <i>ura3</i>	This study
pRS416e-siz1	Derived from pRS416e, with <i>SIZ1</i> gene cassette added	This study
pRS416e-ykl071w	Derived from pRS416e, with <i>YKL071W</i> gene cassette added	This study
pRS416e-zwf1	Derived from pRS416e, with <i>ZWF1</i> gene cassette added	This study
pRS416e-msn2	Derived from pRS416e, with <i>MSN2</i> gene cassette added	This study
pRS416e-ald6	Derived from pRS416e, with <i>ALD6</i> gene cassette added	This study
pRS416e-adh7	Derived from pRS416e, with <i>ADH7</i> gene cassette added	This study
pRS416e-ari1	Derived from pRS416e, with <i>ARI1</i> gene cassette added	This study

Tolerance assay - calculation of maximum specific growth rates

Maximum specific growth rate was used as an indicator for the cellular tolerance towards various inhibitors [12,50,51]. For calculating the maximum specific growth rates of *S. cerevisiae* strains, stationary-phase cells grown in SC medium were transferred into 5 mL of SC medium containing a specific inhibitor (0.8 g/L furfural, 1.2 g/L furfural, 2.0 g/L furfural, 1.26 g/L HMF or 1.72 mg/L menadione) in a 15-mL round-bottom Falcon tube (30°C, 250 rpm). The initial OD₆₀₀ was 0.2. The maximum specific growth rate was determined from the maximum slope of the OD₆₀₀ values over time.

Fermentation

Batch fermentations were carried out as follows: a single colony grown on a YPAD plate was inoculated into 3 mL of SC medium containing 20 g/L glucose in a 15-mL round-bottom Falcon tube and grown until saturation (30°C, 250 rpm). About 400 µL of the stationary-phase cells were transferred into 25 mL of fresh SC media containing 0.8 g/L furfural in 250 mL non-baffled shake flasks. Cells were grown under oxygen-limited conditions (30°C, 100 rpm) as previously reported [46]. The initial OD₆₀₀ was 0.2.

HPLC analysis

The samples were centrifuged and the supernatants were diluted five to ten times before HPLC analysis. An Agilent 1100 series HPLC (Agilent Technologies, Palo Alto, CA, USA) coupled with an Agilent ZORBAX 80A Extend-C18 column was used for detection of furfural and furfuryl alcohol. HPLC parameters were as follows: solvent A, water; solvent B, acetonitrile; 5% B for 15 minutes, then 100% B for 5 minutes, followed by 5% B for 5 minutes; flow rate 1 mL/minute; detection by UV spectroscopy at 277 nm (furfural) or 210 nm (furfuryl alcohol). Under such conditions, furfural and furfuryl alcohol were eluted at 6.7 minutes and 5.6 minutes, respectively. An HPLC system equipped with a refractive index detector (Shimadzu Scientific Instruments, Columbia, MD, USA) was used to

analyze the concentrations of glucose and ethanol in the broth. To separate glucose and ethanol, an HPX-87H column (BioRad, Hercules, CA, USA) was used as described [46].

Additional file

Additional file 1: Figure S1. Sequencing of 17 randomly picked plasmids from the RNAi library. Locations have been mapped to the *S. cerevisiae* genome. Each column represents one chromosome, the height of which is proportional to the size of the indicated chromosome. Each horizontal bar indicates the location of a fragment. **Figure S2.** Sequencing result of pRS416-TTrcx-siz1, which contains a fragment of gene *SIZ1* (underlined). **Figure S3.** Sequencing result of pRS416-TTrcx-gcn4, which contains a fragment of gene *GCN4* (underlined). **Table S1.** Primers used in this study. **Table S2.** Construction of plasmids. **Table S3.** Maximum specific growth rates of strain BAD and its derivatives cultured in SC medium containing 20 g/L glucose. **Table S4.** Maximum specific growth rates of strain BAD and its derivatives cultured in SC medium containing different concentrations of furfural. **Table S5.** Fermentation parameters and estimation of carbon balance in strain BAD and *siz1Δ* after 30 h in SC medium containing 20 g/L glucose and 0.8 g/L furfural.

Abbreviations

5-HMF: 5-hydroxymethylfurfural; OD: optical density; NAPDH: reduced nicotinamide adenine dinucleotide phosphate; RAGE: RNAi-assisted genome evolution; RNAi: RNA interference; ROS: reactive oxygen species; SC: synthetic complete; SC-LEU: synthetic complete medium deficient in leucine; SC-URA: synthetic complete medium deficient in uracil; SUMO: small ubiquitin-like modifier.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All the experiments were performed by HX. Both authors contributed to designing the experiments, writing the manuscript and have approved the final manuscript.

Acknowledgements

The project was supported by the Center for Industrial Biotechnology at the University of Illinois at Urbana-Champaign. We thank Dr William Metcalf (Department of Molecular and Cellular Biology, University of Illinois at Urbana-Champaign, Urbana, USA) for providing *E. coli* strain WM1788, Dr Mingzi M Zhang (Metabolic Engineering Research Laboratory, Institute of Chemical and Engineering Sciences, Agency for Science, Technology and Research, Singapore), Mr Tong Si, Mr Ryan E Cobb, Dr. Jing Liang (Department of Chemical and Biomolecular Engineering, University of Illinois at Urbana-Champaign, Urbana, USA), Dr Xueyang Feng, Mr Jiazhang Lian and Ms Sijin Li (Department of Chemical and Biomolecular Engineering, Institute for Genomic Biology, University of Illinois at Urbana-Champaign) for helpful suggestions and thoughtful discussions.

Received: 28 February 2014 Accepted: 12 May 2014
Published: 23 May 2014

References

- Han K-H, Ko J-H, Yang SH: Optimizing lignocellulosic feedstock for improved biofuel productivity and processing. *Biofuel Bioprod Bior* 2007, **1**:135–146.
- Kumar R, Singh S, Singh OV: Bioconversion of lignocellulosic biomass: biochemical and molecular perspectives. *J Ind Microbiol Biotechnol* 2008, **35**:377–391.
- Pienkos PT, Zhang M: Role of pretreatment and conditioning processes on toxicity of lignocellulosic biomass hydrolysates. *Cellulose* 2009, **16**:743–762.
- Jeong TS, Um BH, Kim JS, Oh KK: Optimizing dilute-acid pretreatment of rapeseed straw for extraction of hemicellulose. *Appl Biochem Biotech* 2010, **161**:22–33.
- Delgenes JP, Moletta R, Navarro JM: Effects of lignocellulose degradation products on ethanol fermentations of glucose and xylose by *Saccharomyces cerevisiae*, *Zymomonas mobilis*, *Pichia stipitis*, and *Candida shehatae*. *Enzyme Microb Technol* 1996, **19**:220–225.
- Miller EN, Jarboe LR, Turner PC, Pharkya P, Yomano LP, York SW, Nunn D, Shanmugam KT, Ingram LO: Furfural inhibits growth by limiting sulfur assimilation in ethanologenic *Escherichia coli* strain LY180. *Appl Environ Microbiol* 2009, **75**:6132–6141.
- Palmqvist E, Grage H, Meinander NQ, Hahn-Hagerdal B: Main and interaction effects of acetic acid, furfural, and p-hydroxybenzoic acid on growth and ethanol productivity of yeasts. *Biotechnol Bioeng* 1999, **63**:46–55.
- Heer D, Sauer U: Identification of furfural as a key toxin in lignocellulosic hydrolysates and evolution of a tolerant yeast strain. *Microb Biotechnol* 2008, **1**:497–506.
- Mussatto SI, Roberto IC: Alternatives for detoxification of diluted-acid lignocellulosic hydrolysates for use in fermentative processes: a review. *Bioresour Technol* 2004, **93**:1–10.
- Allen SA, Clark W, McCaffery JM, Cai Z, Lanctot A, Slininger PJ, Liu ZL, Gorsich SW: Furfural induces reactive oxygen species accumulation and cellular damage in *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 2010, **3**:2.
- Lin FM, Qiao B, Yuan YJ: Comparative proteomic analysis of tolerance and adaptation of ethanologenic *Saccharomyces cerevisiae* to furfural, a lignocellulosic inhibitory compound. *Appl Environ Microbiol* 2009, **75**:3765–3776.
- Park SE, Koo HM, Park YK, Park SM, Park JC, Lee OK, Park YC, Seo JH: Expression of aldehyde dehydrogenase 6 reduces inhibitory effect of furan derivatives on cell growth and ethanol production in *Saccharomyces cerevisiae*. *Bioresour Technol* 2011, **102**:6033–6038.
- Liu ZL: Molecular mechanisms of yeast tolerance and in situ detoxification of lignocellulose hydrolysates. *Appl Microbiol Biotechnol* 2011, **90**:809–825.
- Mills TY, Sandoval NR, Gill RT: Cellulosic hydrolysate toxicity and tolerance mechanisms in *Escherichia coli*. *Biotechnol Biofuels* 2009, **2**:26.
- Ma M, Liu ZL: Comparative transcriptome profiling analyses during the lag phase uncover *YAP1*, *PDR1*, *PDR3*, *RPN4*, and *HSF1* as key regulatory genes in genomic adaptation to the lignocellulose derived inhibitor HMF for *Saccharomyces cerevisiae*. *BMC Genomics* 2010, **11**:660.
- Heer D, Heine D, Sauer U: Resistance of *Saccharomyces cerevisiae* to high concentrations of furfural is based on NADPH-dependent reduction by at least two oxidoreductases. *Appl Environ Microbiol* 2009, **75**:7631–7638.
- Liu ZL, Moon J: A novel NADPH-dependent aldehyde reductase gene from *Saccharomyces cerevisiae* NRRL Y-12632 involved in the detoxification of aldehyde inhibitors derived from lignocellulosic biomass conversion. *Gene* 2009, **446**:1–10.
- Miller EN, Jarboe LR, Yomano LP, York SW, Shanmugam KT, Ingram LO: Silencing of NADPH-dependent oxidoreductase genes (*yqhD* and *dkgA*) in furfural-resistant ethanologenic *Escherichia coli*. *Appl Environ Microbiol* 2009, **75**:4315–4323.
- Si T, Luo Y, Bao Z, Zhao H: RNAi-assisted genome evolution in *Saccharomyces cerevisiae* for complex phenotype engineering. *ACS Syn Bio* 2014. doi:10.1021/sb500074a.
- Liu HP: Constructing yeast libraries. *Method Enzymol* 2002, **350**:72–86.
- Online Sequence Analysis Tools. http://tools.neb.com/~posfai/project/Yeast_Digests.html.
- Statistics of Randomized Library Construction. <http://guinevere.otago.ac.nz/mlrgd/STATS/>.
- Shao Z, Zhao H, Zhao H: DNA assembler, an *in vivo* genetic method for rapid construction of biochemical pathways. *Nucleic Acids Res* 2009, **37**:e16.
- Kebaara BW, Atkin AL: Long 3'-UTRs target wild-type mRNAs for nonsense-mediated mRNA decay in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 2009, **37**:2771–2778.
- Sasano Y, Watanabe D, Ukibe K, Inai T, Ohtsu I, Shimoi H, Takagi H: Overexpression of the yeast transcription activator *MSN2* confers furfural resistance and increases the initial fermentation rate in ethanol production. *J Biosci Bioeng* 2012, **113**:451–455.
- Gorsich SW, Dien BS, Nichols NN, Slininger PJ, Liu ZL, Skory CD: Tolerance to furfural-induced stress is associated with pentose phosphate pathway genes *ZWF1*, *GND1*, *RPE1*, and *TKL1* in *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* 2006, **71**:339–349.
- Miura K, Jin JB, Hasegawa PM: Sumoylation, a post-translational regulatory process in plants. *Curr Opin Plant Biol* 2007, **10**:495–502.
- Geiss-Friedlander R, Melchior F: Concepts in sumoylation: a decade on. *Nat Rev Mol Cell Biol* 2007, **8**:947–956.
- Cheng CH, Lo YH, Liang SS, Ti SC, Lin FM, Yeh CH, Huang HY, Wang TF: SUMO modifications control assembly of synaptonemal complex and polycomplex in meiosis of *Saccharomyces cerevisiae*. *Genes Dev* 2006, **20**:2067–2081.
- Geddes CC, Peterson JJ, Roslander C, Zacchi G, Mullinnix MT, Shanmugam KT, Ingram LO: Optimizing the saccharification of sugar cane bagasse using dilute phosphoric acid followed by fungal cellulases. *Bioresource Technol* 2010, **101**:1851–1857.
- Kim IS, Sohn HY, Jin I: Adaptive stress response to menadione-induced oxidative stress in *Saccharomyces cerevisiae* KNU5377. *J Microbiol* 2011, **49**:816–823.
- Zaldivar J, Martinez A, Ingram LO: Effect of selected aldehydes on the growth and fermentation of ethanologenic *Escherichia coli*. *Biotechnol Bioeng* 1999, **65**:24–33.
- Natarajan K, Meyer MR, Jackson BM, Slade D, Roberts C, Hinnebusch AG, Marton MJ: Transcriptional profiling shows that GCN4p is a master regulator of gene expression during amino acid starvation in yeast. *Mol Cell Biol* 2001, **21**:4347–4368.
- Hinnebusch AG: Mechanisms of gene regulation in the general control of amino acid biosynthesis in *Saccharomyces cerevisiae*. *Microbiol Rev* 1988, **52**:248–273.
- Johnson ES: Protein modification by SUMO. *Annu Rev Biochem* 2004, **73**:355–382.
- Johnson ES, Gupta AA: An E3-like factor that promotes SUMO conjugation to the yeast septins. *Cell* 2001, **106**:735–744.
- Strunnikov AV, Aravind L, Koonin EV: *Saccharomyces cerevisiae* SMT4 encodes an evolutionarily conserved protease with a role in chromosome condensation regulation. *Genetics* 2001, **158**:95–107.
- Li T, Mullen JR, Slagle CE, Brill SJ: Stimulation of *in vitro* sumoylation by SLX5-SLX8: evidence for a functional interaction with the SUMO pathway. *DNA Repair* 2007, **6**:1679–1691.
- Rosonina E, Duncan SM, Manley JL: Sumoylation of transcription factor GCN4 facilitates its SRB10-mediated clearance from promoters in yeast. *Genes Dev* 2012, **26**:350–355.
- Denison C, Rudner AD, Gerber SA, Bakalarski CE, Moazed D, Gygi SP: A proteomic strategy for gaining insights into protein sumoylation in yeast. *Mol Cell Proteomics* 2005, **4**:246–254.
- Srikumar T, Lewicki MC, Raught B: A global *S. cerevisiae* small ubiquitin-related modifier (SUMO) system interactome. *Mol Syst Biol* 2013, **9**:668.
- Treger JM, Schmitt AP, Simon JR, McEntee K: Transcriptional factor mutations reveal regulatory complexities of heat shock and newly identified stress genes in *Saccharomyces cerevisiae*. *J Biol Chem* 1998, **273**:26875–26879.
- Wong CM, Zhou Y, Ng RWM, Kung HF, Jin DY: Cooperation of yeast peroxiredoxins TSA1p and TSA2p in the cellular defense against oxidative and nitrosative stress. *J Biol Chem* 2002, **277**:5385–5394.
- Rep M, Proft M, Remize F, Tamas M, Serrano R, Thevelein JM, Hohmann S: The *Saccharomyces cerevisiae* SKO1p transcription factor mediates HOG pathway-dependent osmotic regulation of a set of genes encoding enzymes implicated in protection from oxidative damage. *Mol Microbiol* 2001, **40**:1067–1083.

45. Izawa S, Maeda K, Miki T, Mano J, Inoue Y, Kimura A: **Importance of glucose-6-phosphate dehydrogenase in the adaptive response to hydrogen peroxide in *Saccharomyces cerevisiae*.** *Biochemical J* 1998, **330**(Pt 2):811–817.
46. Du J, Yuan Y, Si T, Lian J, Zhao H: **Customized optimization of metabolic pathways by combinatorial transcriptional engineering.** *Nucleic Acids Res* 2012, **40**:e142.
47. Radonjic M, Andrau JC, Lijnzaad P, Kemmeren P, Kockelkorn TT, van Leenen D, van Berkum NL, Holstege FC: **Genome-wide analyses reveal RNA polymerase II located upstream of genes poised for rapid response upon *S. cerevisiae* stationary phase exit.** *Mol Cell* 2005, **18**:171–183.
48. Sikorski RS, Hieter P: **A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*.** *Genetics* 1989, **122**:19–27.
49. Gietz RD, Schiestl RH, Willems AR, Woods RA: **Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure.** *Yeast* 1995, **11**:355–360.
50. Kelly C, Jones O, Barnhart C, Lajoie C: **Effect of furfural, vanillin and syringaldehyde on *Candida guilliermondii* growth and xylitol biosynthesis.** *Appl Biochem Biotechnol* 2008, **148**:97–108.
51. Otero JM, Cimini D, Patil KR, Poulsen SG, Olsson L, Nielsen J: **Industrial systems biology of *Saccharomyces cerevisiae* enables novel succinic acid cell factory.** *Plos One* 2013, **8**:e54144.

doi:10.1186/1754-6834-7-78

Cite this article as: Xiao and Zhao: Genome-wide RNAi screen reveals the E3 SUMO-protein ligase gene *SIZ1* as a novel determinant of furfural tolerance in *Saccharomyces cerevisiae*. *Biotechnology for Biofuels* 2014 **7**:78.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

