



Published in final edited form as:

Dev Cell. 2010 December 14; 19(6): 872–883. doi:10.1016/j.devcel.2010.10.023.

Integration of light and brassinosteroid signaling pathways by a GATA transcription factor in *Arabidopsis*

Xiao-Min Luo^{1,5,¶}, Wen-Hui Lin^{1,2,¶}, Shengwei Zhu^{1,2,¶}, Jia-Ying Zhu^{1,5,¶}, Yu Sun², Xi-Ying Fan^{1,5}, Menglin Cheng¹, Yaqi Hao², Eunkyoo Oh², Miaomiao Tian³, Lijing Liu³, Ming Zhang⁴, Qi Xie³, Kang Chong¹, and Zhi-Yong Wang^{2,*}

¹ Key Laboratory of Photosynthesis and Environmental Molecular Biology, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

² Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305

³ Laboratory of Plant Molecular Signaling, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

⁴ GeneExp, San Jose, CA 95112, USA

⁵ Graduate School of the Chinese Academy of Sciences, Beijing 100049, China

SUMMARY

Light and brassinosteroid (BR) antagonistically regulate the developmental switch from etiolation in the dark to photomorphogenesis in the light in plants. Here we identify GATA2 as a key transcriptional regulator that mediates the crosstalk between BR- and light-signaling pathways. Overexpression of GATA2 causes constitutive photomorphogenesis in the dark, whereas suppression of GATA2 reduces photomorphogenesis caused by light, BR deficiency, or the constitutive photomorphogenesis mutant *cop1*. Genome profiling and chromatin immunoprecipitation experiments show that GATA2 directly regulates genes that respond to both light and BR. BR represses GATA2 transcription through the BR-activated transcription factor BZR1, whereas light causes accumulation of GATA2 protein and feedback inhibition of GATA2 transcription. Dark-induced proteasomal degradation of GATA2 is dependent on the COP1 E3 ubiquitin ligase, and COP1 can ubiquitinate GATA2 *in vitro*. This study illustrates a molecular framework for antagonistic regulation of gene expression and seedling photomorphogenesis by BR and light.

INTRODUCTION

Light and brassinosteroid (BR) are key signals that determine the development program of young seedlings. To reach the surface of soil, seedlings that germinate in the dark undergo skotomorphogenesis, exhibiting elongated hypocotyls, small and folded cotyledons with undifferentiated chloroplasts, and repression of light-induced genes. Exposure to light causes a developmental switch from skotomorphogenesis to photomorphogenesis, resulting in short hypocotyls, open and expanded cotyledons, and differentiation of chloroplast (Wei and Deng, 1996). Genetic studies have identified many components that mediate this

*To whom correspondence should be addressed: Z.-Y. W., Tel: 650-325-1521 Ext. 205; Fax: 650-325-6857; zywang24@stanford.edu.

¶These authors contributed equally to the work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

developmental switch by light. Two classes of photoreceptors, phytochrome and cryptochrome, perceiving red/far-red and blue light respectively, play major roles in promoting photomorphogenesis. A group of proteins termed CONSTITUTIVE PHOTOMORPHOGENIC/DE-ETIOLATED/FUSCA (COP/DET/FUS), which are components of the ubiquitination system or COP9 signalosome, are central repressors of photomorphogenesis (Deng et al., 1991; Wei and Deng, 1996). Several classes of transcription factors, such as the b-zip protein LONG HYPOCOTYL 5 (HY5) and the Phytochrome Interacting Factor (PIF) family of b-HLH proteins, directly regulate light-responsive gene expression and are degraded by the ubiquitin system in a light-dependent manner (Chen et al., 2004; Leivar et al., 2008; Ma et al., 2002; von Arnim et al., 1997; Wang et al., 2001). Through these components, light turn on a transcription program that supports photomorphogenic development (Chen et al., 2004; Jiao et al., 2007).

In addition to these light-signaling components, BR also plays a key role in photomorphogenesis. BR deficient mutants show typical de-etiolation phenotypes in the dark with elevated expression of many light-induced genes (Chory et al., 1991; Li et al., 1996; Song et al., 2009; Szekeres et al., 1996). While light inhibits hypocotyl elongation and promotes chlorophyll accumulation, BR promotes hypocotyl elongation and reduces chlorophyll level. BR is perceived by the cell surface receptor kinase BRI1 and downstream signal transduction activates the BZR family transcription factors (Gendron and Wang, 2007), which mediate BR-responsive gene expression (He et al., 2005). Recent studies have established a complete BR signal transduction pathway from the BRI1 to the BZR transcription factors (Kim et al., 2009; Tang et al., 2010; Kim and Wang, 2010). Activation of BZR1 and BZR2 is essential for skotomorphogenesis, as the constitutive photomorphogenesis phenotype of BR-deficient or insensitive mutants are suppressed by the dominant *bzr1-1D* and *bes1-D* mutations, which cause constitutive activation of BR-responsive gene expression (Wang et al., 2002; Yin et al., 2002). It has been proposed that light might inhibit BR synthesis or signaling to inhibit skotomorphogenesis and promote photomorphogenesis (Kang et al., 2001). However, no significant difference in BR level was observed between dark-grown and light-grown plants (Symons et al., 2008). On the other hand, physiological studies of BR-deficient *Arabidopsis* suggested that BR regulates phytochrome- and cryptochrome-mediated responses (Luccioni et al., 2002; Neff et al., 1999). The molecular mechanism of such BR-light interactions has remained unclear.

Analyses of light-responsive promoters have identified a number of light-response promoter elements (LREs), including the G-box, GATA and GT1 motifs (Terzaghi and Cashmore, 1995). It has been suggested that combinations of LREs, rather than individual elements, confer proper light-responsiveness to a promoter (Puente et al., 1996; Terzaghi and Cashmore, 1995). For example, the combination of G-box with GATA element is critical for promoter activation in response to the signals from multiple photoreceptors as well as for repression by the COP/DET system (Chattopadhyay et al., 1998b). Most of the light-signaling transcription factors identified so far bind to the G-box (Liu et al., 2008; Jiao et al., 2007). The transcription factor that regulates light-responsive genes through the essential GATA element has not been identified in plants (Arguello-Astorga and Herrera-Estrella, 1998; Chattopadhyay et al., 1998b; Jiao et al., 2007; Terzaghi and Cashmore, 1995). In fungi, such as *Neurospora*, two GATA-type factors bind to GATA element and regulate gene expression in response to light signal (Scazzocchio, 2000). It has long been proposed that members of the *Arabidopsis* GATA family of transcription factors might play a similar role (Jeong and Shih, 2003; Manfield et al., 2007), however, genetic evidence for this hypothesis is absent.

In this study, we identify a GATA-type transcription factor (GATA2) as a junction between light and BR pathways. Overexpression and loss-of-function experiments demonstrate that

GATA2 is a major positive regulator of photomorphogenesis that mediates a gene expression profile with significant overlap to those caused by light treatment or BR-deficiency. BR-activated BZR1 directly represses GATA2 transcription, whereas light signaling stabilizes the GATA2 protein, likely by inhibiting a COP1-dependent degradation process. The results demonstrate that GATA2 is not only a key light-signaling transcription factor but also a junction for the crosstalk between the BR and light-signaling pathways. The results support a mode of BR-light antagonism through transcriptional and posttranslational regulation of common transcription factors.

RESULTS

GATA2 is a positive regulator of photomorphogenesis

The suppression of the photomorphogenesis phenotype of *bri1* by the *bzr1-1D* mutation suggests that BR inhibits photomorphogenesis through BZR1 and its downstream target genes. Based on BR responsive expression and the presence of BR-response elements in their promoters (He et al., 2005), two BR-repressed genes encoding GATA-type transcription factors, *GATA2* and *GATA4* were considered putative target genes of BZR1. Since previous studies of GATA sequence in light-responsive promoter implicated unknown GATA factors in light-responsive gene expression (Chattopadhyay et al., 1998b), we tested whether GATA2 and GATA4 play a role in light- or BR-regulated gene expression and photomorphogenesis.

GATA2 and GATA4 are two closest members of the subfamily I of *Arabidopsis* GATA factors (Reyes et al., 2004). Quantitative RT-PCR analysis confirmed that the transcript level of *GATA2* is reduced by BR treatment. *GATA2* is expressed at a higher level in the dark than in the light, and BR repression is also more obvious in the dark than in the light (Figure 1A). *GATA2* RNA level is increased in the BR-deficient mutant *det2* and BR-insensitive mutant *bri1-116*, but repressed by the *bzr1-1D* mutation (Figure 1B). A *GATA2* promoter-GUS reporter gene showed strong expression in hypocotyls and petioles (Figure S1), where cell elongation is most sensitive to light and BR. *GATA2* expression was also detected in root tips, the junctions of floral organs, and styles of plants grown under light (Figure S1). RT-PCR assays confirmed ubiquitous expression of *GATA2* in various tissues (Figure S1I). A *GATA2*-YFP fusion protein is localized in the nucleus (Figure S1J–O). Such expression pattern and subcellular localization of *GATA2* is consistent with a role as transcription factor for photomorphogenesis. Recent co-expression analysis has shown that *GATA2* and *GATA4* show strong coexpression with each other (Manfield et al., 2007).

To investigate the function of these GATA factors, we generated transgenic plants over expressing *GATA2* and *GATA4* under the control of the cauliflower mosaic virus 35S promoter (*GATA-ox*). Of five *GATA2-ox* transgenic lines four lines exhibited obvious short hypocotyls and open cotyledons in the dark, similar to the BR-deficient or insensitive mutants (Figure 1C, 1D and S1P–Q). Similarly five of ten *GATA4-ox* lines also showed shorter hypocotyl phenotypes, however the overall phenotypes were weaker than the *GATA2-ox* lines (Figure S1R). We further generated *GATA2* antisense (*GATA-AS*) and artificial microRNA (*GATA-AM*) transgenic plants. The constructs contain conserved sequence and are expected to also suppress *GATA4*. Many *GATA-AS* and *GATA-AM* lines showed long hypocotyl phenotypes in the light (Figure 1E and S2A), but not in the dark (Figure S2G). These results demonstrate that *GATA2* plays an important role in promoting photomorphogenesis, and *GATA4* is likely to play a similar but less prominent role.

As positive regulator of photomorphogenesis, the increased expression of *GATA2* in BR mutants is likely to contribute to the de-etiolation phenotypes. To determine if *GATA2* play a role in BR regulation of photomorphogenesis, we crossed the *GATA-AS* and *GATA-AM*

lines with the BR deficient mutant *det2* and BR insensitive mutant *bin2*, and these plants exhibited longer hypocotyls than the *det2* and *bin2* single mutants (Figure 2A, 2B, S2B, and S2C). We also crossed the *GATA2-ox* line with *bzr1-1D*, which suppresses the de-etiolation phenotypes of the BR-biosynthetic or signaling mutants (Figure 1D). Seedlings homozygous for both *GATA2-ox* and *bzr1-1D* had short hypocotyls and open cotyledon in the dark, resembling the phenotype of *GATA2-ox* (Figure 2C, and S2D), consistent with GATA2 acting downstream of BZR1. These results support an important role of repressing GATA2 in BR inhibition of photomorphogenic development.

Light regulates seedling development through several photoreceptor families that absorb light of distinct wavelengths. To test if GATA2 functions in any specific photoreceptor pathway, we grew the *GATA2-ox* and *GATA*-knockdown lines under monochromatic red, far-red, or blue light. The *GATA2-ox* plants had shorter hypocotyls and the *GATA-AS* or *GATA-AM* plants showed longer hypocotyls under all wavelengths of light but not in the dark (Figure 2D, 2E, and S2E–G), suggesting that GATA2 is likely to function downstream of all photoreceptors. Fluence-rate response analyses indicate that the *GATA2-ox* plants have enhanced sensitivity and the *GATA-AS* and *AM* plants have reduced sensitivity to light (Figure 2E, 2F, and S2G). To test if GATA2 is downstream of the master photomorphogenic repressor COP1 (Deng et al., 1991), we crossed the *GATA-AS* and *GATA-AM* lines into the *cop1-4* and *cop1-6* mutants. Knockdown of GATA partly suppressed the de-etiolation phenotypes of the *cop1* mutants (Figure 2G, S2H), suggesting that GATA2 functions downstream of COP1 in the light signaling pathway.

GATA2 overexpression causes similar transcriptomic changes as light and BR deficiency

To further understand the function of GATA2 in the light- and BR-signaling pathways, we compared the transcriptomic changes caused by *GATA2* overexpression, *bri1* mutation, and light treatment. Four-day old dark-grown seedlings of *GATA2-ox*, *bri1-116*, and wild type were analyzed by microarray using the ATH1 array (Affymetrix). The results showed that expression of 2910 genes were altered in *GATA2-ox* plants, with 1743 genes repressed and 1167 activated by *GATA2* overexpression (>2 fold and $p < 0.05$, Table S1a). In the *bri1-116* mutant, 2992 genes were differentially expressed compared to wild type, and about 38% (1144 of the 2992) of them were also affected in *GATA2-ox* (Figure 3A, Table S1b). More striking overlap was observed for the 120 most-repressed genes in *GATA2-ox*: 103 (86%) of them were also repressed in *bri1-116* (Table S1c). Overall about 93% (1055) of the 1144 co-regulated genes were affected in the same way by *GATA2-ox* and *bri1-116* (Figure 3B, Table S1d). Such similar genomic effects of *GATA2-ox* and *bri1* mutation are consistent with elevated *GATA2* expression in *bri1* contributing to its altered gene expression and de-etiolation phenotype.

When the gene expression changes of *GATA2-ox* were searched against an *Arabidopsis* microarray database that includes 1450 treatments (Zhang et al., 2010), the top nine best matches were microarray experiments that compared seedlings grown under various light conditions to those grown in the dark. The percent overlaps with the light datasets ranged from 27% to 48% (Table S2). The Pearson correlation coefficients of pair-wise comparison between the *GATA2-ox* vs WT data and various light vs dark data range from 0.57 to 0.75 (Table S2), suggesting that *GATA2* overexpression causes a similar genomic response as light exposure. About 47% (1378) of the genes affected in *GATA2-ox* were affected by at least one of the light conditions (Table S3). Among these, 802 genes were affected by *bri1* mutation (Figure 3A, Table S1e). About eighty seven percent of these shared genes were up or down regulated similarly by *GATA2-ox*, the *bri1-116* mutation, and light treatments (Figure 3B, Table S1e). Such similar effects of *GATA2* overexpression, *bri1* mutation, and light on large numbers of genes strongly support an important role for GATA2 in mediating the antagonistic effects of BR and light on gene expression and photomorphogenesis.

GATA2 directly regulates genes that respond to light and BR deficiency

Quantitative RT-PCR assays confirmed that the expression levels of light-repressed genes, such as *TIP2* and *IAA6*, were repressed in *GATA2-ox* and *bri1-116* plants but increased in the *GATA-AS* and *GATA-AM* plants, whereas the levels of light-induced genes, such as *CAB2*, *PSAH2*, were increased in *GATA2-ox* and *bri1-116* plants but reduced in the *GATA-AS* and *GATA-AM* plants (Figure 3C). Chromatin immunoprecipitation (ChIP) assays for GATA sequence-containing regions of promoters demonstrated that GATA2 binds strongly to the promoters of *TIP2*, *CAB2*, *CYCP2.1*, *RBCS1A*, and *PSAH2*, and binds weakly to *IAA6*, *ERD14*, *GA4*, and *FAD5*, which are responsive to light treatment and affected in *bri1* and the *GATA2* transgenic plants. In contrast, GATA2 does not bind to *PSAL*, which is a light-responsive gene not affected by *bri1* or *GATA-AS* (Figure 3D and Table S4). Furthermore, ChIP assays showed GATA2 binding to additional seven genes strongly repressed and three genes strongly activated in *GATA2-ox*, but not to the control gene *UBC30* or two *LHCB* genes that were not affected in *GATA2-ox* (Table S4). These results demonstrate that GATA2 directly activates some of the light-induced and BR-repressed genes and inhibits light-repressed and BR-induced genes.

Light induces accumulation of GATA2 protein, which directly feedback inhibits its own transcription

As a positive regulator of photomorphogenesis, *GATA2* is expected to be activated by light. However, *GATA2* and *GATA4* are expressed at a higher level in dark-grown plants than in light-grown plants (Manfield et al., 2007). Quantitative RT-PCR analysis showed that the transcript levels of *GATA2* and *GATA4* rapidly decreased upon light treatment of dark-grown seedlings (Figure 4A). Interestingly, immunoblot analysis demonstrated that light treatment increased the GATA2 protein accumulation (Figure 4B). Opposite responses at protein and RNA levels are often caused by feedback inhibition of transcription by the protein product of the gene. Indeed, the levels of the endogenous *GATA2* and *GATA4* RNAs were reduced in the *GATA2-ox* transgenic plants, which overexpress the *GATA2* RNA from the transgene (Figure 4C). Overexpression of *GATA2* also led to reduced expression of a *GATA2-GUS* reporter gene in tobacco leaf cell (Figure S3). Chromatin immunoprecipitation assays further showed that the GATA2 protein directly binds to its own promoter *in vivo* (Figure 4D). These results demonstrate that light induces GATA2 protein accumulation at a posttranscriptional level, and light-activated GATA2 protein feedback inhibits the transcription of *GATA2* and *GATA4*.

GATA2 is degraded in the dark by the proteasome in a COP1-dependent manner

Several light-signaling transcription factors, such as HY5 and HFR1, are targeted for proteasomal degradation by the COP1 ubiquitin ligase, and light signaling stabilizes these transcription factors by inactivating COP1 (Osterlund et al., 2000; von Arnim et al., 1997). We investigated whether a similar COP1-dependent process is involved in light regulation of GATA2 accumulation. Treatment of dark-grown seedlings with MG132, an inhibitor of the proteasome, caused GATA2 protein accumulation, indicating that GATA2 is degraded by the proteasome in the dark. Upon transition from light to dark, GATA2 protein level decreased dramatically, and this decrease was blocked by MG132 treatment (Figure 5A), suggesting that light inhibits proteasomal degradation of GATA2. Immunoblotting data showed that the GATA2 protein level was increased in the *cop1* mutants grown in the dark (Figure 5B), indicating that GATA2 degradation requires COP1. The accumulation of GATA2 obviously contributes to the de-etiolation phenotype of *cop1*, as suppressing *GATA2* RNA levels in the *cop1* mutant reduced the GATA2 protein level and increased the hypocotyl length (Figure 2G and Figure 5B). Furthermore, *in vitro* pull-down assays showed that COP1 can directly interact with GATA2 (Figure 5C). *In vitro* ubiquitination assay confirmed that COP1 can ubiquitinate GATA2 *in vitro* (Figure 5D). These results strongly

support the possibility that the GATA2 protein is negatively regulated by COP1-dependent ubiquitination and proteasomal degradation, and inactivation of COP1 by light signaling leads to GATA2 accumulation.

BZR1 binds to the *GATA2* promoter *in vivo*

To test if BZR1 directly regulates *GATA2* expression, we performed chromatin immunoprecipitation followed by quantitative PCR (ChIP-qPCR) assays using *pBZR:BZR1*-CFP transgenic plants and anti-GFP antibody, with 35S-GFP transgenic plants as a control. As shown in Figure 6A, BZR1 bound strongly to the *GATA2* promoter in the dark-grown seedlings but only weakly in the light-grown seedlings, consistent with a prominent role of BR in repressing *GATA2* expression in the dark (Figure 1A). In contrast, BZR1 bound to the *DWF4* promoter strongly in both dark and light conditions. Such transcriptional regulation of *GATA2* leads to altered levels of GATA2 protein, as BR treatment of the *det2* mutant dramatically reduced the GATA2 protein levels (Figure 6B). In contrast, BR and the BR biosynthetic inhibitor brassinazole had little effect on the GATA2 protein level in the transgenic plants that constitutively express *GATA2* from the 35S promoter (Figure S4), suggesting that BR represses *GATA2* at the transcriptional but not posttranscriptional level.

Light does not have a strong effect on BR signaling

Our observation of differential BZR1 binding to the *GATA2* promoter in the dark and light suggests that light affects BZR1 activity. Because BZR1's nuclear localization and DNA binding activity are tightly controlled by BR-regulated phosphorylation (Gendron and Wang, 2007), light could alter BZR1's phosphorylation status if light has an effect on BR level or BR signal transduction. We therefore performed immunoblotting experiments to test whether light affects BZR1 accumulation and phosphorylation (Figure 6C). The results show that plants grown in the dark and under red light or white light conditions contain similar levels of phosphorylated and unphosphorylated BZR1, whereas treatment with BR caused dramatic dephosphorylation of BZR1 (Figure 6C). These results indicate that light does not have a significant effect on BR level or BR signaling upstream of BZR1.

DISCUSSION

Interactions between light and endogenous hormones are critical for plant development. It has been long recognized that BR plays a major role in light regulated plant development. The underlying molecular mechanism has remained unclear. This study identifies members of the GATA factor family (*GATA2* and *GATA4*) as key transcription factors that integrate the BR and light-signaling pathways for coordinated regulation of gene expression and photomorphogenesis (Figure 6D). We show that *GATA2* directly binds to light-responsive promoters *in vivo* and controls the expression of large numbers of genes that respond to both light and BR signaling. *GATA2* is inhibited by BR signaling at the transcriptional level through BZR1 binding to its promoter and is activated by light at the protein level through inhibiting COP1-dependent proteolysis.

GATA2 also binds to its own promoter to feedback inhibit its own transcription. Such feedback mechanism could serve an important desensitizing mechanism during transition from dark to light, but would also lead to de-repression of *GATA2* transcription and incomplete switch to skotomorphogenesis when *GATA2* protein degradation is accelerated in the dark. As such repression of *GATA2* expression by BR becomes essential for maintaining complete skotomorphogenesis in the dark. BR deficiency causes overexpression of *GATA2*, which contributes to de-etiolation in the dark. This study demonstrates a mode of BR-light crosstalk, in which BR signaling inhibits light responses through transcriptional repression of key components of the light-signaling pathway.

GATA2 is a key component for light-responsive gene expression

Analyses of light-regulated promoters have shown an essential role of the GATA element in light-regulated gene expression (Chattopadhyay et al., 1998b; Jeong and Shih, 2003; Terzaghi and Cashmore, 1995). It has been shown that combinations of different LREs, rather than individual elements, confer proper light-responsiveness to a promoter (Puenta et al., 1996). The GATA element functions together with the G-box or GT1 motifs to confer normal response to a wide spectrum of light signals involving multiple photoreceptors and the COP/DET/FUS complex (Chattopadhyay et al., 1998a). These results indicated a role of the GATA element as an essential partner with other LREs in light-regulated gene expression. Previous studies have only identified the G-box-binding factors, including PIFs, HY5, and CIB1. This study identifies GATA2 and GATA4 as the missing transcription factors that act through the GATA element.

Our results provide strong genetic and molecular evidence for the role of GATA2 in light regulation of gene expression and photomorphogenesis. First, overexpression of *GATA2* causes a typical de-etiolation phenotype and a transcriptomic change that resembles those caused by light exposure, whereas suppression of *GATA2* by RNAi or antisense had an opposite effect on hypocotyl elongation and gene expression. Although the long hypocotyl phenotypes of the *GATA-AM* and *-AS* plants are relatively weak, this is likely due to incomplete suppression of *GATA2* expression and/or redundant function of other homologous GATA factors. Second, ChIP assays showed that *in vivo* GATA2 binds to many light-responsive promoters at regions containing GATA motifs, providing direct evidence for GATA2 regulation of light-responsive genes. Finally, GATA2 protein is stabilized by light signaling, most likely through a COP1-dependent mechanism similar to the regulation of the light-signaling transcription factors HY5 and HFR1. GATA2 accumulates in the *cop1* mutants and can interact with COP1 and be ubiquitinated by COP1 *in vitro*, though direct *in vivo* interaction is yet to be demonstrated. Therefore, GATA2 meets the criteria for a primary light signaling transcription factor.

GATA factors are a class of highly conserved transcription factors with a type IV zinc finger followed by a basic region, which are known to recognize the consensus sequence WGATAR (where W is T or A and R is G or A) (Lowry and Atchley, 2000). GATA factors are found in all eukaryotes from fungi to plants and metazoans. In fungi, GATA factors are involved in a number of different processes, ranging from nitrogen utilization, mating-type switch, and light responses (Scazzocchio, 2000). In *Neurospora crassa*, the *White Collar-1* (*WC1*) and *White Collar-2* (*WC2*) loci encoding “plant-like” GATA factors are required for light and circadian responses (Ballario and Macino, 1997). In addition to the GATA DNA binding domain, *WC1* also contains a light-oxygen-voltage (LOV) domain and functions as a photoreceptor (Cheng et al., 2003). It seems that a function of GATA factors in light responses has been conserved during evolution from fungi to higher plants.

The function of *Arabidopsis* GATA factors in light response was not uncovered in previous genetic analysis, and suppression of *GATA2* and *GATA4* only partially suppressed *cop1* and *det2* mutants; these are most likely because of genetic redundancy. The *Arabidopsis* genome contains 29 genes that encode GATA factors (Reyes et al., 2004). Some members of the GATA family have been shown to play a role in regulating flower development (Zhao et al., 2004), chlorophyll synthesis, and carbon/nitrogen metabolism (Bi et al., 2005; Mara and Irish, 2008). *In vitro* DNA binding assays have shown binding of GATA1 to the *GATA* elements of the *GAPB* promoter that are essential for light responsive expression (Jeong and Shih, 2003). GATA2 shares 76% sequence identity with GATA1 in the DNA binding domain and is likely to have similar DNA binding specificity for GATA elements. Our ChIP experiment shows that GATA2 binds to promoter regions containing GATA sequence.

Additional GATA family members may be involved in light responses, as their expression levels are regulated by light. Higher expression in the light-grown than dark-grown seedlings has been observed for *GATA6*, *GATA7*, *GATA21*, *GATA22*, and *GATA23* (Manfield et al., 2007). None of these genes are affected in the *GATA2-ox* plants based on our microarray data, suggesting their light regulation is independent of *GATA2*. In fact *GATA21* and *GATA22* are induced by red light in a PIF3-dependent manner (Monte et al., 2004). In contrast, four other genes, *GATA2*, *GATA4*, *GATA9*, and *GATA12*, showed stronger expression in the dark-grown than light-grown seedlings (Manfield et al., 2007), and they are all repressed in the *GATA2-ox* plants. Based on similarity in sequence, gene structure, and expression profiles, these four *GATA* genes have been predicted to share common ancestry, with *GATA2* and *GATA4* arisen from a recent chromosomal duplication (Reyes et al., 2004). *GATA2* and *GATA4* are coexpressed with each other and share common coexpressed genes, which include *PHYA* and light-signaling transcription factors *PIF3*, *PIF1/PIL5*, and *HFR1* (Manfield et al., 2007), consistent with their role in light signaling. In contrast, *GATA9* and *GATA12* do not show significant coexpression with any of the genes known to be involved in light signaling. It has been suggested that *GATA9* and *GATA12* have diverged from *GATA2* and *GATA4* in expression and possibly in function as well (Manfield et al., 2007). Based on our expression microarray data, only *GATA2* and *GATA4*, but not *GATA9* and *GATA12*, are overexpressed in the *bri1* mutant more than two fold and repressed in the *bri1 bri1-ID* double mutant, indicating that BR regulates the transcription of *GATA2* and *GATA4* but not their close homologs *GATA9* and *GATA12*. Further genetic analysis of double or multiple loss-of-function mutants will be required to understand whether other GATA factors also play a role in photomorphogenesis.

The relationship between *GATA2* and other light-signaling transcription factors is key for understanding light-responsive gene expression. Several lines of evidence suggest that *GATA2* functions together with the G-box-binding factors. First, *GATA* and G-box elements are found together in many light responsive promoters, and their dual presence is essential for normal light responsiveness in a synthetic promoter (Chattopadhyay et al., 1998b). Second, *GATA2* shows strong coexpression with *PIF3*, *PIF1/PIL5*, *SPT*, and *HFR1* (Manfield et al., 2007), many of which bind to the G-box. Third, *GATA2* is stabilized by light at the posttranslational level, likely through the same COP1-dependent mechanism that regulates *HY5* and *HFR1*. It is also worth noting that a higher percentage of the genes up regulated than down regulated in *GATA2-ox* are *HY5* targets (Lee et al., 2007) (27% of 1167 up-regulated genes vs 21% of 1743 down-regulated genes), which is consistent with our hypothesis that *GATA* and G-box elements together confer light activated expression by recruiting *GATA2* and *HY5*. Whether *GATA2* directly interacts with other light-signaling transcription factors and how they orchestrate dynamic light-regulated gene expression are yet to be analyzed in future studies.

GATA2 is a key junction for the antagonism between BR and light signaling pathways

Genetic studies have long demonstrated a critical role of BR in skotomorphogenesis (Li et al., 1996; Szekeres et al., 1996). The antagonizing relationship between BR and light has been analyzed at the genetic and physiological levels. Mutations that reduce BR level enhanced the light responses (Neff et al., 1999), and a rice *phyB* mutant showed enhanced BR responses (Jeong et al., 2007). The antagonism at the level of gene expression was recognized in the initial studies of the BR deficient mutants (Chory et al., 1991) (Li et al., 1996; Szekeres et al., 1996), and confirmed by our microarray data showing similar transcriptomic changes caused by the *bri1* mutation and light exposure. The similar effects of BR deficiency and light on seedling development and expression of large numbers of genes suggested three possible mechanisms of interaction between the BR and light-signaling pathways: (1) light reduces BR level or BR sensitivity, (2) BR regulates light-

signaling components to inhibit light signaling, or (3) BR and light signaling pathways regulate common target genes through separate transcription factors independently controlled by each pathway. This study provides evidence for the second mechanism of BR-light crosstalk, and recent genomic analysis of BZR1 target genes supported the presence of also the third mechanism (Sun et al., 2010).

A previous study proposed that light inhibits BR biosynthesis by repressing a small G protein that binds to and activates a BR-biosynthetic enzyme (Kang et al., 2001). However, subsequent direct BR measurement failed to detect significant difference in BR levels between light-grown and dark-grown plants, but showed light reducing the level of gibberellin, another hormone that also promotes cell elongation (Symons and Reid, 2003). Our observations of no obvious effect of light on the phosphorylation status and accumulation of BZR1 or on BZR1 binding to the *DWF4* promoter are consistent with the lack of change of BR level by light. Our results further suggest that light does not inhibit BR signaling upstream of BZR1. However, stronger BZR1 binding to the *GATA2* promoter was observed in the dark-grown than light-grown seedlings. It is possible that light has an effect on the availability of BZR1 binding site or BZR1-interacting proteins at the *GATA2* promoter. In contrast to the lack of strong effect of light on BR signaling, BR obviously has a strong effect on light signaling by repressing *GATA2* expression.

Our results show that *GATA2* plays a key role in BR regulation of photomorphogenesis. *GATA2* accumulates in the *det2* mutant and *GATA2* knockdown partially suppresses the photomorphogenic phenotypes of dark-grown *det2* and *bin2*, indicating that de-etiolation in the BR mutants are at least partly due to the increased levels of *GATA2*. About one third of the genes affected in *bri1* are affected similarly by *GATA2-ox*, suggesting that the elevated *GATA2* level contributes to a major portion of *bri1*'s effect on genome expression and that BZR1 repression of *GATA2* is a major mechanism for BR inhibition of light responses. By inhibiting transcription and promoting protein accumulation of *GATA2*, respectively, BR and light antagonistically regulate the level of *GATA2* activity and consequently the expression of its downstream target genes. Thus, *GATA2* represents a key junction of crosstalk between BR and light signaling pathways.

The mechanism of BR-light crosstalk through *GATA2* is distinct from those for light crosstalk with GA and cytokinin. In addition to light repression of GA level, GA also affects the activity or accumulation of the light-signaling transcription factors PIF/PIL and HY5 (Alabadi et al., 2008; de Lucas et al., 2008; Feng et al., 2008). The DELLA proteins of the GA signaling pathway directly interact with and inhibit members of the PIF/PIL family, which are negative regulators of photomorphogenesis (de Lucas et al., 2008; Feng et al., 2008). GA also promotes degradation of HY5, possibly through a COP1-dependent process (Alabadi et al., 2008). In contrast, cytokinin, which promotes photomorphogenesis, induces HY5 protein accumulation (Vandenbussche et al., 2007). Whether other hormones also regulate *GATA2* to modulate light responses remains to be tested by future studies.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

The wild type, various mutants, and transgenic *Arabidopsis thaliana* plants were in the *Columbia* ecotype. Seeds were sterilized by incubation in freshly prepared 10% bleach plus 0.01% Triton X-100 for 15 min and then washed 3–4 times with sterilized water. The surface-sterilized seeds were treated in 4 °C for 2 days and at 22 °C under white light for 8 hours to induce uniform germination. For phenotype analyses, seedlings were grown on 0.8% phytoagar plates containing half-strength Murashige-Skoog (MS) nutrient and 1% sucrose. White light (about 100 $\mu\text{mol}/\text{m}^2/\text{s}$) was provided by fluorescence light source in a

growth room at 22 °C. Growth under red, far-red, and blue light was carried out in a LED light chamber (E-30LEDL3, Percival) at 22 °C. Seedlings were photographed next to a size reference (ruler) and their hypocotyl lengths measured using the Image J software. Seeds were harvested from plants grown in a greenhouse supplemented to 16-hour light/day and a temperature range of 18–28 °C.

Vector construction and transformation

A 1152-bp genomic fragment containing full-length *GATA2* open reading frame was amplified by PCR and then cloned into the BamHI and KpnI sites of the pSN1301 binary vector to place *GATA2* under the control of the CaMV 35S promoter.

The *GATA2* antisense construct was made by inserting the *GATA2* full-length cDNA fragment in reverse orientation into the pSN1301 plasmid. The artificial microRNA constructs were made using the vectors and methods previously reported (Schwab et al., 2006) (See Supplemental Information for details). The 35S::*GATA2-YFP* fusion construct was generated by inserting a full-length *GATA2* cDNA without stop codon fused to the N terminus of the pEZR-LNY vector.

The *GATA2-ox*, *GATA2-AS*, *GATA2-AM* and 35S::*GATA2-YFP* binary constructs were transformed into the *Agrobacterium tumefaciens* strain GV3101 and then introduced into *Arabidopsis thaliana* Columbia wild-type plants via a floral dip method. About 20 T1 transgenic lines with single T-DNA insertion were selected for further analysis. Homozygous T3 or T4 transgenic seedlings were used for phenotype and molecular characterization.

Protein expression and antibody preparation

The full length *GATA2* cDNA was cloned into the pGEX-4T-1 vector to express GST-*GATA2* protein in *E. coli* Rosetta cells (Novagen). The recombinant fusion protein was purified using glutathione-agarose beads (GE Healthcare) and used to immunize rabbit. The anti-*GATA2* antibody was purified from the immune serum using immobilized GST-*GATA2* (Aminolink® Immobilization Kit, Pierce Biotechnology). The anti-Histone H3 antibody for loading control was from Millipore (Catalogue No. 07-690)

Total RNA isolation and quantitative RT-PCR analysis

Total RNA was extracted from *Arabidopsis* seedlings using the Trizol RNA extraction kit (Invitrogen, USA). The first-strand cDNA was synthesized by using M-MLV reverse transcriptase (Promega, USA) and used as RT-PCR templates. Quantitative real-time PCR analyses were carried out on Mx3000P (Stratagene, USA) by using the SYBR® Green reagent (TOYOBO, JAPAN) according to the manufacturer's instructions. The RT-PCR was repeated at least three times using samples harvested separately. The *UBC30* gene was used as internal reference. See Supplemental Information for primer sequences used for RT-PCR.

Microarray Data Analysis

Arabidopsis seedlings (Columbia, *GATA2-ox*, *bri1-116*) were grown on 1/2 MS medium in the dark for 4.5 days, and the seedlings were frozen in liquid nitrogen in complete darkness, and then the *bri1-116* seedlings were selected from the segregating population. Ten micrograms (10 µg) of total RNA from the seedlings was used to prepare probes for hybridization, and each probe was hybridized independently to one chip according to the protocol of the ATH1 array manufacturer (Affymetrix). Three independent biological repeats were conducted. The data were analyzed using Genespring software ver. 7. Data that were flagged as absent, using the Affymetrix mismatch probes, in two or more of the repeats for each genotype were removed. Genes that passed this filter for any one of the genetic

backgrounds were used for further analysis. P-value <0.05 and fold change >2 (for *GATA2-ox*) or fold change >1.8 (for *bri1-116*) were used to identify genes differentially expressed in *GATA2-ox* or *bri1-116* compared to wild type control seedlings.

To determine what experimental conditions causes similar gene expression changes as *GATA2-ox*, we carried out expression fingerprint searching by comparing the differential gene expression pattern between *GATA2-ox* treatment and all available 1450 treatment/control microarray comparisons (T/Cs) in the Gene Expression Browser (GEB) database (<http://www.expressionbrowser.com/>) (Zhang et al., 2010). We inputted the pairs of *GATA2-ox* significant (2-fold and P<0.05 as cutoff) gene IDs and their log₂ ratios, and compared them to each T/C of GEB with the following procedure: (1) Select the significant genes from the T/C using 2-fold and P<0.05 as cutoff. (2) Compute the overlapping genes between *GATA2-ox* and the T/C. The chi-square test was used for filtering out non-significant overlaps (P<0.01 as cutoff). (3) Compute the Pearson's correlation coefficient using the paired log₂ ratios of *GATA2-ox* and the T/C for the overlapping genes. The significance of correlation P value was also computed to reject non-significant correlations (p<0.01 as cutoff). As a result, all hits were significant in both number of overlapping genes and expression changes (Pearson correlation). Finally, the hit list was ordered by the Pearson correlation coefficient.

Chromatin immunoprecipitation

Chromatin immunoprecipitation experiments were performed following the protocol described previously (He et al., 2005), using 2-week-old light-grown wild type and *35S::GATA2-YFP* transgenic *Arabidopsis* seedlings or 5-day-old dark- and light-grown *35S-GFP* and *pBZR::BZR1-CFP* seedlings. An affinity-purified anti-GFP polyclonal antibody was used to immunoprecipitate the BZR1 or GATA2 protein-DNA complex, and the precipitated DNA was analyzed by real-time PCR using the SYBR[®] Green reagent (TOYOBO, JAPAN). Results were presented as the ratio of the amount of DNA immunoprecipitated from BZR1-CFP or GATA2-YFP samples to that of the control samples (*35S-GFP* or wide type). The *UBC30* and *PP2A* genes were used as the negative controls. The CHIP experiments were performed 3 times, from which the means and standard deviations were calculated. The primer sequences for CHIP-qPCR are in Supplemental Information.

Protein purification and pull down assay

The GST-GATA2 protein was expressed using the pGEX-4T-1 vector in *E. coli* Rosetta cells (Novagen). The recombinant fusion protein was purified using glutathione-agarose beads (GE Healthcare). For pull-down assay, COP1 fused to maltose binding protein (MBP) was purified using amylose resin (NEB). Glutathione beads containing GST-GATA2 was incubated with MBP, MBP-COP1. The mixture was rotated in a cold room for 1 hr and the beads were washed 5 times with wash buffer (20 mM Tris-HCl [pH8.0], 200 mM NaCl). The proteins were eluted from the beads by boiling in equal volume of 2×SDS buffer and loaded onto a SDS-PAGE gel. Gel blots were analyzed using an anti-MBP antibody (NEB).

In vitro ubiquitination assay

The MBP-COP1 and GST-GATA2 proteins expressed in *E. coli* were affinity purified for *in vitro* ubiquitination assays. To improve the E3 activity of MBP-COP1, the purified MBP-COP1 and MBP control proteins on maltose beads were incubated with *Arabidopsis* cell extract for 30 min. After incubation, the cell extract was removed and the beads were washed. To perform the *in vitro* ubiquitination assay, crude extract containing recombinant wheat E1 (GI: 136632), human E2 (UBCh5b), His-UBI (UBQ14), purified GST-GATA2 and purified MBP-COP1 (or MBP control) were incubated at 30 °C with agitation in an

Eppendorf Thermomixer for 1.5 hr. The proteins were immunoblotted after SDS-PAGE and GST-GATA2 was detected using an anti-GST antibody.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by grants from the National Institute of General Medical Sciences (R01GM066258), National Science Foundation of China (90817009), and Ministry of Agriculture of China (2008ZX08009-003). We thank Dr. Xing Wang Deng for providing COP1 clones and *cop1* mutants, and Dr. Winslow Briggs for comments on the manuscript.

REFERENCE

- Alabadi D, Gallego-Bartolome J, Orlando L, Garcia-Carcel L, Rubio V, Martinez C, Frigerio M, Iglesias-Pedraz JM, Espinosa A, Deng XW, et al. Gibberellins modulate light signaling pathways to prevent *Arabidopsis* seedling de-etiolation in darkness. *Plant J.* 2008; 53:324–335. [PubMed: 18053005]
- Arguello-Astorga G, Herrera-Estrella L. Evolution of Light-Regulated Plant Promoters. *Annu Rev Plant Physiol Plant Mol Biol.* 1998; 49:525–555. [PubMed: 15012245]
- Ballario P, Macino G. White collar proteins: PASsing the light signal in *Neurospora crassa*. *Trends Microbiol.* 1997; 5:458–462. [PubMed: 9402704]
- Bi YM, Zhang Y, Signorelli T, Zhao R, Zhu T, Rothstein S. Genetic analysis of *Arabidopsis* GATA transcription factor gene family reveals a nitrate-inducible member important for chlorophyll synthesis and glucose sensitivity. *Plant J.* 2005; 44:680–692. [PubMed: 16262716]
- Chattopadhyay S, Ang LH, Puente P, Deng XW, Wei N. *Arabidopsis* bZIP protein HY5 directly interacts with light-responsive promoters in mediating light control of gene expression. *Plant Cell.* 1998a; 10:673–683. [PubMed: 9596629]
- Chattopadhyay S, Puente P, Deng XW, Wei N. Combinatorial interaction of light-responsive elements plays a critical role in determining the response characteristics of light-regulated promoters in *Arabidopsis*. *Plant J.* 1998b; 15:69–77. [PubMed: 9744096]
- Chen M, Chory J, Fankhauser C. Light signal transduction in higher plants. *Annu Rev Genet.* 2004; 38:87–117. [PubMed: 15568973]
- Cheng P, He Q, Yang Y, Wang L, Liu Y. Functional conservation of light, oxygen, or voltage domains in light sensing. *Proc Natl Acad Sci U S A.* 2003; 100:5938–5943. [PubMed: 12719523]
- Chory J, Nagpal P, Peto CA. Phenotypic and Genetic Analysis of *det2*, a New Mutant That Affects Light-Regulated Seedling Development in *Arabidopsis*. *Plant Cell.* 1991; 3:445–459. [PubMed: 12324600]
- de Lucas M, Daviere JM, Rodriguez-Falcon M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blazquez MA, Titarenko E, Prat S. A molecular framework for light and gibberellin control of cell elongation. *Nature.* 2008; 451:480–484. [PubMed: 18216857]
- Deng XW, Caspar T, Quail PH. *cop1*: a regulatory locus involved in light-controlled development and gene expression in *Arabidopsis*. *Genes Dev.* 1991; 5:1172–1182. [PubMed: 2065972]
- Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, Chen L, Yu L, Iglesias-Pedraz JM, Kircher S, et al. Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature.* 2008; 451:475–479. [PubMed: 18216856]
- Gendron JM, Wang ZY. Multiple mechanisms modulate brassinosteroid signaling. *Curr Opin Plant Biol.* 2007; 10:436–441. [PubMed: 17904409]
- He JX, MJ, Gendron, Sun YL, SS, Gampala, Gendron N, Sun CQ, Wang ZY. BZR1 is a transcriptional repressor with dual roles in brassinosteroid homeostasis and growth responses. *Science.* 2005; 307:1634–1638. [PubMed: 15681342]
- Jeong DH, Lee S, Kim SL, Hwang I, An G. Regulation of brassinosteroid responses by phytochrome B in rice. *Plant Cell Environ.* 2007; 30:590–599. [PubMed: 17407537]

- Jeong MJ, Shih MC. Interaction of a GATA factor with cis-acting elements involved in light regulation of nuclear genes encoding chloroplast glyceraldehyde-3-phosphate dehydrogenase in *Arabidopsis*. *Biochem Biophys Res Commun*. 2003; 300:555–562. [PubMed: 12504119]
- Jiao Y, Lau OS, Deng XW. Light-regulated transcriptional networks in higher plants. *Nat Rev Genet*. 2007; 8:217–230. [PubMed: 17304247]
- Kang JG, Yun J, Kim DH, Chung KS, Fujioka S, Kim JI, Dae HW, Yoshida S, Takatsuto S, Song PS, et al. Light and brassinosteroid signals are integrated via a dark-induced small G protein in etiolated seedling growth. *Cell*. 2001; 105:625–636. [PubMed: 11389832]
- Kim TW, Guan S, Sun Y, Deng Z, Tang W, Shang J, Sun Y, Burlingame AL, Wang ZY. Brassinosteroid signal transduction from cell-surface receptor kinases to nuclear transcription factors. *Nat Cell Bio*. 2009; 11:1254–1260. [PubMed: 19734888]
- Kim TW, Wang ZY. Brassinosteroid Signal Transduction from Receptor Kinases to Transcription Factors. *Annu Rev Plant Biol*. 2010; 61:681–704. [PubMed: 20192752]
- Lee J, He K, Stolc V, Lee H, Figueroa P, Gao Y, Tongprasit W, Zhao H, Lee I, Deng XW. Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell*. 2007; 19:731–749. [PubMed: 17337630]
- Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH. Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr Biol*. 2008; 18:1815–1823. [PubMed: 19062289]
- Li J, Nagpal P, Vitart V, McMorris TC, Chory J. A role for brassinosteroids in light-dependent development of *Arabidopsis*. *Science*. 1996; 272:398–401. [PubMed: 8602526]
- Liu H, Yu X, Li K, Klejnot J, Yang H, Lisiero D, Lin C. Photoexcited CRY2 interacts with CIB1 to regulate transcription and floral initiation in *Arabidopsis*. *Science*. 2008; 322:1535–1539. [PubMed: 18988809]
- Lowry JA, Atchley WR. Molecular evolution of the GATA family of transcription factors: conservation within the DNA-binding domain. *J Mol Evol*. 2000; 50:103–115. [PubMed: 10684344]
- Luccioni LG, Oliverio KA, Yanovsky MJ, Boccalandro HE, Casal JJ. Brassinosteroid mutants uncover fine tuning of phytochrome signaling. *Plant Physiol*. 2002; 128:173–181. [PubMed: 11788763]
- Ma L, Gao Y, Qu L, Chen Z, Li J, Zhao H, Deng XW. Genomic evidence for COP1 as a repressor of light-regulated gene expression and development in *Arabidopsis*. *Plant Cell*. 2002; 14:2383–2398. [PubMed: 12368493]
- Manfield IW, Devlin PF, Jen CH, Westhead DR, Gilmartin PM. Conservation, convergence, and divergence of light-responsive, circadian-regulated, and tissue-specific expression patterns during evolution of the *Arabidopsis* GATA gene family. *Plant Physiol*. 2007; 143:941–958. [PubMed: 17208962]
- Mara CD, Irish VF. Two GATA transcription factors are downstream effectors of floral homeotic gene action in *Arabidopsis*. *Plant Physiol*. 2008; 147:707–718. [PubMed: 18417639]
- Monte E, Tepperman JM, Al-Sady B, Kaczorowski KA, Alonso JM, Ecker JR, Li X, Zhang Y, Quail PH. The phytochrome-interacting transcription factor, PIF3, acts early, selectively, and positively in light-induced chloroplast development. *Proc Natl Acad Sci U S A*. 2004; 101:16091–16098. [PubMed: 15505214]
- Neff MM, Nguyen SM, Malanchruvil EJ, Fujioka S, Noguchi T, Seto H, Tsubuki M, Honda T, Takatsuto S, Yoshida S, et al. BAS1: A gene regulating brassinosteroid levels and light responsiveness in *Arabidopsis*. *Proc Natl Acad Sci U S A*. 1999; 96:15316–15323. [PubMed: 10611382]
- Osterlund MT, Wei N, Deng XW. The roles of photoreceptor systems and the COP1-targeted destabilization of HY5 in light control of *Arabidopsis* seedling development. *Plant Physiol*. 2000; 124:1520–1524. [PubMed: 11115869]
- Puente P, Wei N, Deng XW. Combinatorial interplay of promoter elements constitutes the minimal determinants for light and developmental control of gene expression in *Arabidopsis*. *EMBO J*. 1996; 15:3732–3743. [PubMed: 8670877]
- Reyes JC, Muro-Pastor MI, Florencio FJ. The GATA family of transcription factors in *Arabidopsis* and rice. *Plant Physiol*. 2004; 134:1718–1732. [PubMed: 15084732]

- Scazzocchio C. The fungal GATA factors. *Curr Opin Microbiol.* 2000; 3:126–131. [PubMed: 10745000]
- Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D. Highly specific gene silencing by artificial microRNAs in *Arabidopsis*. *Plant Cell.* 2006; 18:1121–1133. [PubMed: 16531494]
- Song L, Zhou XY, Li L, Xue LJ, Yang X, Xue HW. Genome-wide analysis revealed the complex regulatory network of brassinosteroid effects in photomorphogenesis. *Mol Plant.* 2009; 2:755–772. [PubMed: 19825654]
- Sun Y, Fan X-Y, Cao D-M, He K, Tang W, Zhu J-Y, He J-X, Bai M-Y, Zhu S, Oh E, et al. Integration of Brassinosteroid Signal Transduction with the Transcription Network for Plant Growth Regulation in *Arabidopsis*. *Dev Cell.* 2010 In Press.
- Symons GM, Reid JB. Hormone levels and response during de-etiolation in pea. *Planta.* 2003; 216:422–431. [PubMed: 12520333]
- Symons GM, Smith JJ, Nomura T, Davies NW, Yokota T, Reid JB. The hormonal regulation of de-etiolation. *Planta.* 2008; 227:1115–1125. [PubMed: 18214530]
- Szekeres M, Nemeth K, Koncz-Kalman Z, Mathur J, Kauschmann A, Altmann T, Redei GP, Nagy F, Schell J, Koncz C. Brassinosteroids rescue the deficiency of CYP90, a cytochrome P450, controlling cell elongation and de-etiolation in *Arabidopsis*. *Cell.* 1996; 85:171–182. [PubMed: 8612270]
- Tang W, Deng Z, Wang ZY. Proteomics shed light on the brassinosteroid signaling mechanisms. *Curr Opin Plant Biol.* 2010; 23:27–33. [PubMed: 20004136]
- Terzaghi WB, Cashmore AR. Light-regulated Transcription. *Annu Rev Plant Physiol Plant Mol Biol.* 1995; 46:445–474.
- Vandenbussche F, Habricot Y, Condiff AS, Maldiney R, Van der Straeten D, Ahmad M. HY5 is a point of convergence between cryptochrome and cytokinin signalling pathways in *Arabidopsis thaliana*. *Plant J.* 2007; 49:428–441. [PubMed: 17217468]
- von Arnim AG, Osterlund MT, Kwok SF, Deng XW. Genetic and developmental control of nuclear accumulation of COP1, a repressor of photomorphogenesis in *Arabidopsis*. *Plant Physiol.* 1997; 114:779–788. [PubMed: 9232869]
- Wang H, Ma LG, Li JM, Zhao HY, Deng XW. Direct interaction of *Arabidopsis* cryptochromes with COP1 in light control development. *Science.* 2001; 294:154–158. [PubMed: 11509693]
- Wang ZY, Nakano T, Gendron J, He J, Chen M, Vafeados D, Yang Y, Fujioka S, Yoshida S, Asami T, et al. Nuclear-localized BZR1 mediates brassinosteroid-induced growth and feedback suppression of brassinosteroid biosynthesis. *Dev Cell.* 2002; 2:505–513. [PubMed: 11970900]
- Wei N, Deng XW. The role of the COP/DET/FUS genes in light control of *Arabidopsis* seedling development. *Plant Physiol.* 1996; 112:871–878. [PubMed: 8938399]
- Yin Y, Wang ZY, Mora-Garcia S, Li J, Yoshida S, Asami T, Chory J. BES1 accumulates in the nucleus in response to brassinosteroids to regulate gene expression and promote stem elongation. *Cell.* 2002; 109:181–191. [PubMed: 12007405]
- Zhang M, Zhang Y, Liu L, Yu L, Tsang S, Tan J, Yao W, Kang MS, An Y, Fan X. Gene Expression Browser: large-scale and cross-experiment microarray data integration, management, search & visualization. *BMC Bioinformatics.* 2010; 11:433. [PubMed: 20727159]
- Zhao Y, Medrano L, Ohashi K, Fletcher JC, Yu H, Sakai H, Meyerowitz EM. HANABA TARANU is a GATA transcription factor that regulates shoot apical meristem and flower development in *Arabidopsis*. *Plant Cell.* 2004; 16:2586–2600. [PubMed: 15367721]




Figure 1. GATA2 is a positive regulator of photomorphogenesis

(A), BR treatment reduces *GATA2* RNA level. *Arabidopsis* seedlings grown in the dark (WD) or light (WL) for five days were treated with mock solution or 100 nM 24-epibrassinolide (BR) for 3 hours and the expression of *GATA2* was analyzed by qRT-PCR. (B) qRT-PCR analysis of *GATA2* and *GATA4* RNA levels in 5-day-old dark-grown wild type (Col-0), *det2*, *bri1-116*, and *bri1-116 bzr1-1D*. (C) Dark-grown phenotypes of three *GATA-ox* lines. Lower panels show qRT-PCR of *GATA2* expression (see also Figure S1P, S1R). (D) Phenotypes of light-grown (first on left) or dark-grown seedlings of wild type (Col-0), BR mutants and a representative *GATA2-ox* transgenic line 6. (E) Phenotypes of antisense (AS) or artificial-microRNA (AM) transgenic *Arabidopsis* seedlings with reduced levels of *GATA2* and *GATA4* (see also Figure S2A for quantitation data). Lower panel shows qRT-PCR analysis of *GATA2* and *GATA4* in these transgenic seedlings. All error bars are standard deviation (SD).




Figure 2. GATA2 acts downstream of both BR and light signaling pathways to promote photomorphogenesis

(A) Phenotypes of *det2* mutants crossed with the *GATA2/4* antisense (AS) or artificial microRNA (AM) lines (see also Figure S2B). (B) Phenotypes of the *bin2* mutant crossed with the *GATA-AS* and *-AM* lines (see also Figure S2C). (C) Phenotypes of *bzi1-1D* mutants crossed with *GATA2-ox* (see also Figure S2D). (D) *GATA2-ox* plants (right of each pair) have short hypocotyls than wild type (left) when grown under red ($26 \mu\text{mol}/\text{m}^2/\text{s}$), blue ($13 \mu\text{mol}/\text{m}^2/\text{s}$) and far-red ($100 \mu\text{mol}/\text{m}^2/\text{s}$) light conditions (see also Figure S2E). (E) Relative hypocotyl lengths of *GATA2-ox* seedlings (L3 line) grown under various fluence rate of red light. (F) Fluence-rate response curve of hypocotyl lengths of *GATA2-AS* and *-AM* lines grown in the dark or various intensities of blue light. Error bars in E and F are SD and significant differences from WT are marked (** $p < 0.01$, * $p < 0.05$). (G) Phenotypes of dark-grown *cop1* mutants crossed with *GATA-AS* or *GATA-AM* lines (see also quantitation data in Figure S2I). The seedlings were grown for 7 days.




Figure 3. GATA2 directly regulates genes that are responsive to both BR and light
 (A) Venn diagrams of the number of genes differentially expressed in the dark-grown *GATA2-ox* vs WT, genes affected in the *bri1-116* mutant, and genes affected in at least one of the light-treatment microarray experiments (see also Table S1). The numbers in the overlapping areas indicate the number of shared genes. (B) Scatter plot of log₂ fold change values of *GATA2-ox* vs WT and *bri1-116* vs WT for 802 genes differentially expressed in dark-grown *GATA2-ox* vs WT, *bri1-116* vs WT, and light-grown vs dark-grown wild type seedlings. Effects of light treatment on the expression are denoted by color as shown (see also Table S1). (C) Quantitative RT-PCR analysis of a number of known light-responsive genes in *GATA2-ox*, *GATA-AS-1*, *GATA-AM-4*, or *bri1-116* plants grown in the dark for 5 days, compared to wild type plants grown in the dark and then untreated or treated with white light for 2 hours. (D) Chromatin immunoprecipitation followed by real time PCR (ChIP-qPCR) assays of GATA2 binding to promoters of genes in panel C, performed using *35S::GATA2-YFP* transgenic and wild type control seedlings grown in light for 2 weeks and an anti-GFP antibody. GATA2 binding was measured by qPCR as the ratio between GATA2-YFP and control sample. The *UBC30* gene was used as a negative control. Error bars indicate SD. (See also Table S4).




Figure 4. Light regulates GATA2 accumulation at the post-translational level

(A) Light represses *GATA2* and *GATA4* transcription levels. Dark-grown *Arabidopsis* seedlings were treated with white light for indicated time and RNA levels of *GATA2* and *GATA4* were measured by real-time qRT-PCR. Error bars indicate standard deviation. (B) Light promotes GATA2 protein accumulation. Immunoblot analysis of GATA2 protein in 5-day-old dark-grown *GATA2-ox* L6 line seedlings treated with white light for the indicated time. (C) qRT-PCR analysis of the levels of RNA expressed from the endogenous *GATA2* and *GATA4* genes (endo) or total GATA2 RNA level in wild type (WT) and the *GATA2-ox* transgenic seedlings (L3 and L6). *UBC30* was used as internal control. (D) ChIP-qPCR analysis of GATA2 binding to its own promoter. The upper panel shows a diagram of the promoter (open box), 5'UTR (black line) and the first exon (black box) of the *GATA2* gene. Black circles indicate positions of putative GATA motifs. Lines marked a to f show GATA2-binding (solid) and non-binding (dashed) regions analyzed by qPCR. The lower panel shows ChIP-qPCR data. Error bars indicate SD.




Figure 5. Light regulates GATA2 accumulation through a COP1 ubiquitin ligase-dependent process

(A) Immunoblot analysis of GATA2 protein levels. Dark-grown (D) or light-grown (L or L/D) 5-day-old *35S:GATA2* transgenic seedlings (L3 line) were treated with mock solution (-) or 10 μ M MG132 (+) for 4 hours in the dark (D and L/D) or light (L). Histone H3 was probed as a loading control. (B) Immunoblot assay of GATA2 protein level in 5-day old dark-grown wild type, the *cop1* mutants, and *cop1-4* crossed with the *GATA-AS* line. (C) *In vitro* pull-down assay showing the interaction between GATA2 and COP1. (D) *In vitro* ubiquitination assay showing ubiquitination of GST-GATA2 by MBP-COP1. The arrow points to the GST-GATA2 band and the star marks the ubiquitinated GST-GATA2 bands.




Figure 6. BR represses *GATA2* transcriptional level through *BZR1* direct binding to its promoter (A) ChIP-qPCR assays of *BZR1* binding to the *GATA2* promoter. The *pBZR1::BZR1-CFP* and *35S::GFP* transgenic *Arabidopsis* seedlings grown in dark or light for 5 days were used in ChIP using anti-GFP antibody. The upper panel shows a diagram depicting the putative promoter (open box), 5'UTR (black line) and the first exon (black box) of the *GATA2* gene. Open and black circles indicate the positions of putative E-box and BRRE motifs, respectively. Thin lines marked P1 to P6 show *BZR1*-binding (solid) and nonbinding (dashed) regions analyzed by qPCR. The lower panel shows the quantitative PCR data for enrichment as ratio between *BZR1*-CFP and *35S*-GFP normalized to the *CNX5* control gene. Error bars indicate SD. (B) Immunoblot analysis shows BR repression of *GATA2*

accumulation. The *det2* seedlings were grown in the dark on medium with 100 nM 24-epibrassinolide (eBL) for 5 days, or grown without eBL for 5 days and then treated with 10 μ M eBL for 0 to 4 hours. The level of Histone H3 was used as a loading control. (C) Light does not have a significant effect on BZR1 phosphorylation status. Phosphorylated (pBZR1) and unphosphorylated (BZR1) BZR1 were analyzed by immunoblotting using an anti-BZR1 antibody in *Arabidopsis* seedlings grown in dark (D), under red (R) or white (W) light for three (3 d) or five days (5 d). Seedlings grown in white light for five days were treated with 100 nM brassinolide (+BL) for 30 min. The gel blot was stained with Ponceau S to show protein loading (the Rubisco major band is weaker in dark-grown samples). (D) A model for GATA2 function in BR- and light-regulation of photomorphogenesis. In the dark, the BR-activated BZR1 directly represses *GATA2* transcription and COP1 promotes *GATA2* ubiquitination and degradation, ensuring a low *GATA2* level for etiolation/skotomorphogenesis. In the presence of light, COP1 is inactivated and the *GATA2* protein accumulates to a high level to promote photomorphogenesis through binding to target genes. The *GATA2* protein also feedback inhibits its own transcription by directly binding to its promoter, potentially desensitizing the system upon light-induced accumulation of *GATA2* protein. When BR levels are low, reduced BZR1 activity leads to overexpression of *GATA2*, which promotes photomorphogenesis.