

Quorum sensing and virulence regulation in *Xanthomonas campestris*

Ya-Wen He & Lian-Hui Zhang

Institute of Molecular and Cell Biology, Singapore, Singapore

Correspondence: Lian-Hui Zhang, Institute of Molecular and Cell Biology, 61 Biopolis Drive, Singapore 138673, Singapore. Tel.: +65 6586 9686; fax: +65 6779 1117; e-mail: lianhui@imcb.a-star.edu.sg

Received 31 January 2008; revised 19 April 2008; accepted 30 April 2008.
First published online 16 June 2008.

DOI:10.1111/j.1574-6976.2008.00120.x

Editor: Miguel Camara

Keywords

DSF; cell–cell communication; c-di-GMP; two-component system; xanthan.

Abstract

It is now clear that cell–cell communication, often referred to as quorum sensing (QS), is the norm in the prokaryotic kingdom and this community-wide genetic regulatory mechanism has been adopted for regulation of many important biological functions. Since the 1980s, several types of QS signals have been identified, which are associated commonly with different types of QS mechanisms. Among them, the diffusible signal factor (DSF)-dependent QS system, originally discovered from bacterial pathogen *Xanthomonas campestris* pv. *campestris*, is a relatively new regulatory mechanism. The rapid research progress over the last few years has identified the chemical structure of the QS signal DSF, established the DSF regulon, and unveiled the general signaling pathways and mechanisms. Particular noteworthy are that DSF biosynthesis is modulated by a novel posttranslational autoinduction mechanism involving protein–protein interaction between the DSF synthase RpfF and the sensor RpfC, and that QS signal sensing is coupled to intracellular regulatory networks through a second messenger cyclic-di-GMP and a global regulator Clp. Genomic and genetic analyses show that the DSF QS-signaling pathway regulates diverse biological functions including virulence, biofilm dispersal, and ecological competence. Moreover, evidence is emerging that the DSF QS system is conserved in a range of plant and human bacterial pathogens.

Introduction

The *Xanthomonas* genus is one of the most ubiquitous groups of plant-associated bacterial pathogens. Members of this genus have been shown to infect at least 124 monocotyledonous and 268 dicotyledonous plant species (Leysn *et al.*, 1984). Among them, *Xanthomonas campestris* is the most dominant species with at least 141 pathovars, which infect a wide range of plants including many of agricultural interest, e.g. cabbage, cauliflower, broccoli (*X. c.* pv. *campestris*), tomato and pepper (*X. c.* pv. *vesicatoria*), cotton (*X. c.* pv. *malvacearum*), soybean (*X. c.* pv. *glycines*), and walnut (*X. c.* pv. *juglandis*) (Hayward, 1993).

Xanthomonas campestris pv. *campestris* (Pammel) Dowson (*Xcc* hereafter) is the causal agent of black rot of crucifers, which is possibly the most important disease of crucifers worldwide (Williams, 1980). *Xcc* is a small, rod-shaped, aerobic gram-negative, nonspore-forming bacterium (Onsando, 1992). The bacterium has a single polar flagellum, is positive in catalase activity and hydrogen

sulfide reaction, oxidase negative, and does not produce nitrate or indole (Onsando, 1992). According to the Commonwealth Mycological Institute (CMI) distribution map, the disease has been recorded in over 90 countries representing all the five continents (Anon, 1978). The most important hosts of *Xcc* are the members of *Brassica oleracea*, including cabbage, cauliflower, broccoli, Brussel sprouts, and kale. The bacterial cells infect cabbage through hydathodes at the leaf margins, causing V-shaped lesions, or through stomata, causing round lesions (Lopes & Quezado-Soares, 1997). Another study shows that *Xcc* normally gains entry into plants via hydathodes (Hugouvieux *et al.*, 1998). Once inside the plant, *Xcc* colonizes the vascular system where it produces an extracellular polysaccharide (EPS) called xanthan, which can obstruct the xylem vessels, causing tissue necrosis and severe leaf wilting (Williams, 1980; Onsando, 1992). Evidence is accumulating that xanthan may also play other roles in infection. Treatment of plants with xanthan suppresses the accumulation of callose and enhances plant susceptibility to *Xcc* (Yun *et al.*, 2006).

Xanthan is also associated with the formation of biofilms, which was proposed to play a protective role for the bacterial cells inside the biofilm structure against plant defense responses (Torres *et al.*, 2007). In addition, *Xcc* is known to encode the biosynthesis of other virulence determinants, including extracellular enzymes, such as endoglucanase and polygalacturonase, and a type III secretion system (also known as the Hrp secretion system), which contribute to the pathogenicity of *Xcc* by providing nutrients through degradation of the xylem walls and facilitating the passage of bacteria via degradation of the bordered pit membranes in the vascular system, or through interference with plant defenses, respectively (Onsando, 1992; Dow & Daniels, 1994). Furthermore, *Xcc* produces a yellow pigment xanthomonadin, which appears to contribute to the bacterial epiphytic survival (Poplawsky *et al.*, 2000).

In addition to its roles in *Xcc* virulence, xanthan has had valuable industry and biotechnology applications. It was known long time ago that *Xcc* is capable of producing substantial quantities of xanthan during the fermentation of carbohydrate substrates (Lilly *et al.*, 1958). Xanthan is a heteropolysaccharide with a primary structure consisting of repeated pentasaccharide units formed by two glucose units, two mannose units, and one glucuronic acid unit (García-Ochoa *et al.*, 2000). Xanthan has distinctive rheological properties in aqueous solutions (Morris *et al.*, 1977). Xanthan is nontoxic, nonsensitizing, and does not cause skin or eye irritation (Kennedy & Bradshaw, 1984). The gum also has unusual stability to temperature, pH, shear, enzyme degradation, and excellent compatibility with salts over a wide pH range (Leela & Sharma, 2000). Owing to its unique properties, xanthan has utility in a wide range of applications. In the food industry, it has been used in a wide variety of foods as a suspending, stabilizing, and thickening agent (Leela & Sharma, 2000). Because of its properties in thickening aqueous solutions, as a dispersing agent, and stabilizer of emulsions and suspensions, xanthan is used in pharmaceutical formulations, cosmetics, and agricultural products. It is used in textile printing pastes, glazes, slurry explosive formulations, and rust removers. Furthermore, the high viscosity of solutions and the water solubility of the polymer have also created important applications for xanthan in the petroleum industry where it is commonly used in drilling fluids and in enhancing oil recovery processes (García-Ochoa *et al.*, 2000). The widespread utility of xanthan has made it an important industrial biopolymer and its global demand continues to increase each year (Sutherland, 1998).

Research progress in the last few years shows that *Xcc* has evolved a unique quorum-sensing (QS) system, which plays a key role in the regulation of xanthan (EPS) biosynthesis and bacterial virulence. The QS system of *Xcc* differs from other known QS systems in various aspects, including the

signal chemistry, autoregulation of signal production, and ways to couple to the bacterial central regulatory networks. This review will focus on these fascinating features, as well as the latest findings on QS regulon and signaling networks. The potential implications of these findings on other bacterial pathogens will also be discussed.

***Xcc* communicates using a new type of language**

It is increasingly evident that bacteria are not isolated solitary organisms, but actively 'speak' to one another using small chemical signaling molecules as languages. QS is one of the sophisticated mechanisms of cell-cell communications, by which bacteria count their own number by producing, detecting, and responding to the accumulation of signaling molecules that they export into their environment (for reviews, see Whitehead *et al.*, 2001; Fuqua & Greenberg, 2002; Bassler & Losick, 2006; Williams *et al.*, 2007). Booming research on QS since the 1980s has led to identification of several types of bacterial QS signals (Fig. 1). Among them, *N*-acyl homoserine lactones (AHLs) seem to be one family of the most widely conserved QS signals. The chemical structure of the first AHL signal from marine bacterium *Vibrio fischeri* was unveiled more than two decades ago (Eberhard *et al.*, 1981). Up till now, AHL-type QS signals, which share a conserved homoserine lactone ring linked to a fatty acid side chain with variations in chain length and substitutions (Fig. 1), have been found in more than 70 bacterial species (for a review, see Williams *et al.*, 2007). Autoinducer 2 (AI-2) is another type of widely detected QS signal (Xavier & Bassler, 2003). At present, two chemically similar AI-2 signals have been identified in *Vibrio harveyi* and *Salmonella typhimurium*, respectively. In *V. harveyi*, AI-2 is (2*S*, 4*S*)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran-borate (Chen *et al.*, 2002); in *S. typhimurium*, AI-2 is (2*R*,4*S*)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran (Miller *et al.*, 2004). The LuxS protein is required for the production of AI-2 (Xavier & Bassler, 2003) and database search shows that the *luxS* gene is widespread and presently found in over 60 species, including members of the *Beta*-, *Gamma*-, and *Delta*- and *Epsilonproteobacteria*, *Spirochaetales*, *Firmicutes*, and *Actinobacteridae*, as well as genera belonging to the *Deinococcus* and *Cytophaga* groups, and the green sulfur bacteria (for reviews, see Vendeville *et al.*, 2005; Williams *et al.*, 2007). Evidence is accumulating that these QS signals play vital roles in coordinating various aspects of bacterial activities, including production of virulence factors and production of biofilms (for reviews, see Zhang & Dong, 2004; Bassler & Losick, 2006; Dong *et al.*, 2007; Williams *et al.*, 2007).

Similar to many other bacterial pathogens, *Xcc* has also evolved QS systems for genetic regulation at the community







Signal	Structure	Representative organism
Gram-negative		
N-acyl homoserine lactone (AHL)		<i>V. fischeri</i> (n=1; R= H) <i>A. tumefaciens</i> (n=2; R=O) <i>E. carotovora</i> (n=2; R=OH) <i>P. aeruginosa</i> (n=4; R=O) <i>V. harveyi</i> (n=0; R=OH)
AI-2 family		<i>V. harveyi</i> (A) <i>S. typhimurium</i> (B)
Hydroxy-palmitic acid methyl ester (PAME)		<i>R. solanacearum</i>
<i>Pseudomonas</i> quinolone signal (PQS)		<i>P. aeruginosa</i>
DSF family		<i>X. campestris</i> <i>S. maltophilia</i> <i>B. cenocepacia</i> <i>X. fastidiosa</i>
Gram-positive		
γ -Butyrolactones (A-factor)		<i>S. griseus</i>

Fig. 1. Chemical structures of representative microbial QS signals. The information was summarized from the following references: Bassler & Losick (2006); Dong *et al.* (2007); Boon *et al.* (2008); Colnaghi Simionato *et al.* (2007); and Huang & Wong (2007a).

level. An early transposon mutagenesis analysis unveiled that a cluster of *Xcc* genes, designated *rpfABCDEFG* (for regulation of pathogenicity factors), are involved in the production of EPS and extracellular enzymes (Tang *et al.*, 1991). Subsequently, it was found that the protease production in the mutant of the *rpfF* gene, which encodes a putative enoyl CoA hydratase, was restored by cocultivation in proximity to *Xcc* wild-type strains, suggesting that *Xcc* wild-type strains could produce a diffusible signal factor (DSF) (Barber *et al.*, 1997). To detect DSF, the original bioassay is based on the ability of DSF to restore endoglucanase production in the *rpfF* mutant (Barber *et al.*, 1997). The detection system was later improved by fusing the promoter of an endoglucanase gene to the β -glucuronidase (GUS) reporter gene (Slater *et al.*, 2000; Wang *et al.*, 2004). This improved DSF sensor, and the finding that mutation of the *rpfC* gene that encodes a hybrid sensor implicated in DSF signaling increases DSF production up to 16-fold (Slater *et al.*, 2000; Wang *et al.*, 2004), have greatly facilitated the purification and characterization of the DSF

signal as *cis*-11-methyl-2-dodecenoic acid (Fig. 1) (Wang *et al.*, 2004).

Comparison of the biological activities of a range of DSF derivatives has led to identification of a few key structural features that determine the signaling activity. The most important feature is the unsaturated double bond at the α,β position of DSF; the saturated derivative is about 20 000 times less active than DSF (Wang *et al.*, 2004). The other features, including the *cis* configuration of the α,β double bond, the chain length, and the methyl substitution at the C-11 position, also contribute to the biological activity of DSF (Wang *et al.*, 2004). Interestingly, two groups of Gram-negative bacterial QS signals, i.e., AHLs produced by many Gram-negative bacterial species and the 3-hydroxypalmitic acid methyl ester from *Ralstonia solanacearum* (Flavier *et al.*, 1997), are also derived from fatty acids but do not contain an α,β unsaturated bond at the fatty acid moieties (Fig. 1). The chemical structures of another two groups of QS signals produced by Gram-negative bacteria, i.e., PQS from *Pseudomonas aeruginosa* and AI-2 from *V. harveyi* and

S. typhimurium, are clearly different from DSF (Fig. 1). The lipopeptide-type QS signal A factor produced by gram-positive *Streptomyces* is also structurally different from DSF (Fig. 1). Collectively, these data indicate that DSF represents a new family of bacterial QS signals.

However, DSF may not be the sole QS-signaling molecule used by *Xcc*. A diffusible factor (DF) is implicated in the regulation of the production of xanthomonadins and EPS in *Xcc* (Poplawsky & Chun, 1997; Poplawsky *et al.*, 1998). The nonpigmented *pigB* mutants of *Xcc* cannot synthesize DF, make fourfold less EPS, cause fewer lesions on cabbage, and are impaired in epiphytic survival (Poplawsky & Chun, 1998). The *Xcc pigB* mutant can be restored extracellularly for xanthomonadin and EPS production by a DF-producing strain, or via addition of partially purified exogenous DF. Similarly, application of the DF extracts on cauliflower plants restores the virulence of the DF-deficient mutant to a level comparable to its wild-type *Xcc* parental strain (Chun *et al.*, 1997). A subsequent study has shown that *pigB* and DF are needed for epiphytic survival because normal xanthomonadin production is critical for protection of the bacterial pathogen against UV light (Poplawsky *et al.*, 2000). MS analysis of DF predicted a butyrolactone structure (Chun *et al.*, 1997), which remains to be validated by nuclear magnetic resonance (NMR) analysis or synthetic chemistry. Similar to DSF, DF is also likely a conserved signal. Most *Xanthomonas* species and some *Streptomyces* strains produce DF (Chun *et al.*, 1997; Poplawsky *et al.*, 2005). Further analysis of the *pigB* DNA sequence reveals the presence of two genes and only one of them (*Xcc4014*), which encodes a pteridine-dependent deoxygenase-like protein, is needed for production of the DF signal (Poplawsky *et al.*, 2005). DF and DSF are clearly distinct signals because they differ in biological activity, chromatographic mobility, chemical structure, and the genes encoding biosynthesis (Barber *et al.*, 1997; Poplawsky *et al.*, 1998, 2005; Wang *et al.*, 2004). The fact that EPS production is influenced by both DF and DSF suggests that the two QS systems may interact with each other at certain regulatory junctions. Utilization of two QS signal molecules to regulate different sets of virulence genes may provide *Xcc* the plasticity in response to different environments. Further characterization of the chemical structure of DF, and its signaling mechanisms in regulation of xanthomonadin and EPS production will be essential for a clear understanding of the roles of the two sophisticated QS systems (DSF and DF) in modulation of *Xcc* physiology and virulence.

DSF regulon and DSF-dependent biological functions

DSF-dependent biological functions were initially determined from the phenotype analysis of the DSF-deficient

mutant *rpfF::Tn5lac* of *Xcc* strain 8004 and *rpfF* deletion mutant $\Delta rpfF$ of *Xcc* strain XC1. Null mutation of *rpfF* in both strains leads to reduced production of EPS and extracellular enzymes, including proteases and cellulases (Tang *et al.*, 1991; Barber *et al.*, 1997; He *et al.*, 2006b). Protease production of the *rpfF* mutant strain (*rpfF::Tn5lac*) could be restored by cultivation in proximity to wild-type strains (Barber *et al.*, 1997). DSF activity could be extracted with ethyl acetate from supernatants of overnight cultures of *Xcc*, and addition of the extracted DSF could restore extracellular enzyme production of the DSF-deficient mutants (Barber *et al.*, 1997; Wang *et al.*, 2004). Similarly, addition of purified or synthetic DSF at a physiologically relevant concentration (1 μ M) to the *rpfF* deletion mutants of strain XC1 restores the production of EPS and extracellular enzymes (He *et al.*, 2006b), suggesting that DSF positively regulates the production of EPS and extracellular enzymes. Another important DSF-dependent biological function in *Xcc* is biofilm formation. Mutation of the *rpfF* gene in both *Xcc* strains 8004 and XC1 causes the formation of bacterial cell aggregates, but whose size varies depending on the strains (Dow *et al.*, 2003; He *et al.*, 2006b). Addition of DSF to the bacterial cultures stops formation of cell aggregates by the *rpfF* deletion mutants, suggesting a critical role of DSF signal in modulation of *Xcc* switching between planktonic and biofilm growth forms.

The availability of the *Xcc* genome sequence and advancement of microarray technology have greatly facilitated the determination of the scope of genes and biological functions regulated by the DSF signal. Genome annotation of *Xcc* strains ATCC33913 and 8004 leads to prediction of 4181 and 4273 genes, respectively (da Silva *et al.*, 2002; Qian *et al.*, 2005). The two *Xcc* genomes are highly similar, sharing over 92% identical or almost identical genes with merely about 2% strain-specific genes (Qian *et al.*, 2005). Measurement of gene expression levels has been performed using an oligonucleotide microarray based on the genome sequence of *Xcc* strain ATCC33913 (He *et al.*, 2006b). Comparison of the global gene expression profiles of wild-type strain XC1 and its DSF-deficient mutant XC1 $\Delta rpfF$ at three growth stages (middle exponential, late exponential, and early stationary phase) identifies 183 genes showing significant changes in expression level (greater than and equal to twofold), whereas altered expression of 421 genes is evident when comparison was made between XC1 $\Delta rpfF$ with and without DSF addition of the DSF signal (He *et al.*, 2006b). Further comparison of the two sets of data establishes an overlapping set of 165 DSF-dependent genes, known as the 'core' members of the DSF regulon (He *et al.*, 2006b). Among them, > 80% are DSF-activated, and < 20% are DSF-repressed.

Interestingly, the DSF-dependent genes are in general randomly distributed throughout the bacterial genome (He *et al.*, 2006b). The pattern is similar to the *P. aeruginosa*

genes regulated by AHL-type QS signals, which are also randomly scattered in the bacterial genome (Schuster *et al.*, 2003). These data suggest that both AHL-type and DSF-type QS systems are unlikely to have been acquired recently by these microorganisms.

Based on the published gene list of *Xcc* strain ATCC33913 (da Silva *et al.*, 2002), except for the 10 genes encoding hypothetical proteins, the products of the remaining 155 DSF-regulated genes are grouped into 12 functional categories: (1) extracellular enzymes, (2) lipopolysaccharide and EPS synthesis and secretion, (3) multidrug resistance and detoxification, (4) flagellum biosynthesis, motility, and chemotaxis, (5) hypersensitive response and pathogenicity (Hrp) system, (6) iron uptake, (7) protein metabolism, (8) tricarboxylic acid (TCA) cycle, (9) aerobic and anaerobic respiration, (10) transcription regulators, (11) membrane components and transporters, and (12) fatty acid metabolism and others (He *et al.*, 2006b). Thus, in addition to these previously known DSF-dependent activities such as production of EPS and extracellular enzymes (Barber *et al.*, 1997), microarray analysis unveils several new functions mediated by DSF, such as flagellum synthesis, resistance to toxin and oxidative stress, iron uptake, and aerobic respiration. Some of these newly identified DSF-mediated phenotypes, including resistance to toxin acriflavin and hydrogen peroxide and bacterial survival at different temperatures, have been verified by genetic and phenotype analysis (He *et al.*, 2006b). Taken together, the available genetic and genomic data suggest that the DSF-signaling system is not only essential for coordinating the expression of virulence genes at the community level but also appears to be of critical importance for maintenance of *Xcc* ecological competence.

Intriguingly, in contrast to the positive-regulation pattern of the DSF signal on most virulence genes, microarray analysis reveals that in YEB medium mutation of *rpfF* encoding DSF biosynthesis significantly increases the expression of the genes encoding the type III secretion system (T3SS, also known as Hrp) (He *et al.*, 2006b). However, the other report shows that when grown in MME medium, mutation of *rpfG* encoding DSF signal transduction has no effect on the expression of *hrpX* and *hrpG*, which encode the master regulators of the *hrp* operon (Ryan *et al.*, 2007). While the impact of DSF on the T3SS remains to be further verified by genetic and genomic analysis under different conditions favorable to the expression of type III secretion genes, it is interesting to note that the AHL-type QS system in *P. aeruginosa* also negatively controls the expression of the T3SS regulon (Blevess *et al.*, 2005). Given that T3SS is one set of key bacterial virulence determinants in many bacterial pathogens (Hueck, 1998; Ghosh, 2004), thorough investigation of the regulatory mechanisms of T3SS by QS and other signaling mechanisms would be of key importance for

designing and developing effective new treatments against bacterial virulence and infections.

DSF-signaling network in *Xcc*

How is the DSF signal detected, transduced, and connected to the intracellular regulatory networks? This intriguing puzzle has been resolved recently by the collective works of several laboratories. Although many details remain to be added on, the general outline of the DSF-signaling network has been established, which could be a useful platform for further in-depth investigations.

RpfC and RpfG constitute a two-component system to sense and to transduce the DSF signal

In the *Xcc* genome, *rpfG*, *rpfH*, and *rpfC* are transcribed as the same operon although *rpfC* has its own weak promoter, suggesting that the three genes could be functionally related (Slater *et al.*, 2000). RpfC is a hybrid sensor kinase consisting of five transmembrane domains (TM), a histidine kinase (HK) domain, a receiver (REC) domain, and a histidine phosphotransfer (HPT) domain. RpfG protein contains two major domains: a receiver domain and an HD-GYP domain with conserved HD and GYP motifs (Fig. 2b). Deletion of either *rpfC* or *rpfG* decreases the production of EPS and extracellular enzymes (Slater *et al.*, 2000), which is reminiscent of the DSF-deficient mutants (Barber *et al.*, 1997; He *et al.*, 2006b). Based on these results, it was proposed that the RpfC/RpfG is a signal transduction system that couples the synthesis of pathogenic factors to sensing of environmental signals that may include DSF itself (Slater *et al.*, 2000).

This speculation was later confirmed by site-directed mutagenesis and domain deletion analysis. Peptide alignment analysis with other well-characterized two-component sensors and regulators reveals four conserved residues, i.e., His¹⁹⁸ in the HK domain, Asp⁵¹² in the receiver domain, His⁶⁵⁷ in the HPT domain of RpfC, and Asp⁸⁰ in the receiver domain of RpfG (He, 2006; He *et al.*, 2006a). Substitution of these residues with unrelated amino acids or deletion of any one of the domains containing these conserved residues results in decreased virulence factor production (He, 2006; He *et al.*, 2006a), suggesting that RpfC/RpfG use the conserved phosphorelay mechanisms to transduce the DSF signal for modulation of virulence factor production. In addition, while exogenous addition of DSF to the DSF-deficient mutant Δ rpfF restores the production of EPS and extracellular enzymes to the wild-type levels, the same treatment has no effect on mutant Δ rpfG and the double-deletion mutant Δ rpfF Δ rpfC (He *et al.*, 2006b). Consistent with the above genetic analysis data, microarray analysis shows that RpfG and RpfC regulons are highly similar to the DSF regulon (He, 2006; He *et al.*, 2006b). Cumulatively,




Fig. 2. Schematic representation of a model of the DSF signal auto-induction. At a low cell density, The DSF sensor RpfC forms a complex with the DSF synthase RpfF through its receiver domain, which prevents the enzyme from effective synthesis of the DSF signal. The dashed arrow indicates basal signal generation or no signal flow. At a high cell density, accumulated extracellular DSF signal interacts with RpfC and induces a conformational change in the sensor, which undergoes autophosphorylation and facilitates release of RpfF and phosphorelay from the sensor to its response regulator RpfG. The event boosts DSF biosynthesis and induces or changes the expression pattern of the DSF regulon. The solid arrow indicates signal flow or signal generation. The question mark indicates unknown precursor(s).

these findings suggest that RpfC and RpfG constitute a two-component system essential for sensing and transducing the DSF signal (Fig. 2b).

DSF signaling is mediated by the second messenger cyclic-di-GMP

Different from the orthodox response regulator that typically contains a receiver domain and a DNA-binding domain (for a review, see Stock *et al.*, 2000), RpfG contains a receiver domain and an HD domain (Slater *et al.*, 2000). Proteins containing the HD domain comprise a superfamily of metal-dependent phosphohydrolases (Aravind & Koonin, 1998). RpfG belongs to a subgroup of the HD superfamily that contains an additional GYP signature motif (Slater

et al., 2000). *In trans* expression of the HD-GYP domain of RpfG alone in the null mutants of RpfG restores the production of EPS and extracellular enzymes to a level close to the wild-type strain, whereas expression of the receiver domain alone has no effect (He, 2006; Ryan *et al.*, 2006), indicating that the regulatory activity of RpfG is associated with the HD-GYP domain. Bioinformatic analysis reveals that in general the number of proteins with the HD-GYP domain in bacterial genomes is positively proportional to the number of GGDEF domain proteins that are implicated in synthesis of cyclic-di-GMP, and is negatively proportional to that of EAL domain proteins, which are implicated in the degradation of cyclic-di-GMP (Römling *et al.*, 2005). These findings, together with the observation revealing that the proteins with HD domains share the conserved motifs with eukaryotic cyclic-nucleotide phosphodiesterases (Aravind & Koonin, 1998), led to an insightful prediction that the proteins containing HD-GYP domains may have phosphodiesterase activity and could be involved in cyclic diguanylate signaling (Galperin *et al.*, 1999).

A recent study presents several lines of evidence that demonstrate that the HD-GYP domain of RpfG is a cyclic-di-GMP phosphodiesterase (Ryan *et al.*, 2006). Firstly, the purified HD-GYP domain protein is able to degrade the model substrate bis(*p*-nitrophenyl) phosphate but has no activity against *p*-nitrophenyl phosphate, indicating that it is a phosphodiesterase but not phosphomonoesterase. Secondly, the purified HD-GYP domain can degrade cyclic-di-GMP to generate GMP. Thirdly, in contrast to the effect on cyclic-di-GMP, the protein has no detectable activity against ATP, GTP, GMP, cGMP, or cAMP. Fourthly, substitution of the H and D residues of the HD-GYP domain abrogates both the enzymatic activity against cyclic-di-GMP and the regulatory activity on virulence factor production. These biochemical and genetic data, together with the findings that the HD-GYP domain alone can substitute RpfG in regulation of EPS and extracellular enzymes production (He, 2006; Ryan *et al.*, 2006), provide a clear indication that the regulatory activity of the response regulator RpfG depends on its enzymatic activity against cyclic-di-GMP, which is a widely conserved novel second messenger implicated in regulation of a wide range of bacterial functions (Römling *et al.*, 2005).

A global transcriptional regulator Clp is involved in DSF signaling

The implication of cyclic-di-GMP in the DSF signaling process and that RpfG acts by degrading cyclic-di-GMP to produce cGMP suggest a possibility that there may be a nucleotide receptor(s) in the downstream signaling pathway. Data mining of the genes influenced by DSF shows that the *clp* gene, which encodes a transcriptional regulator with

a nucleotide-binding domain and a DNA-binding domain, is likely to be a potential candidate (He *et al.*, 2007). The *clp* (CAP-like protein) gene was named due to the observation that its product shares a high similarity to the catabolite activation factor (CAP, also widely known as Crp derived from its function as a cAMP receptor protein) of *Escherichia coli* (de Crecy-Lagard *et al.*, 1990). However, the null mutation of *clp* in *Xcc* strain NRRLB1459 does not affect the utilization of various carbon sources but instead decreases the biosynthesis of EPS, extracellular cellulase, and polygalacturonate lyase (de Crecy-Lagard *et al.*, 1990), the phenotypes similar to those of the DSF-deficient mutants (Barber *et al.*, 1997; Slater *et al.*, 2000; Wang *et al.*, 2004). In addition, the transcriptional expression of several DSF-regulated genes (He *et al.*, 2006a), including the cellulase gene *engXCA*, the protease gene *ptr1*, the flagellin gene *fliC*, and the heat-shock protein gene *groESL*, is also positively regulated by Clp in different *Xcc* strains (de Crecy-Lagard *et al.*, 1990; Hsiao & Tseng, 2002; Lee *et al.*, 2003; Chang *et al.*, 2005; Hsiao *et al.*, 2005). These original findings, as well as the observation that the transcriptional expression of *clp* is influenced by DSF and dependent on the RpfC/RpfG two-component system (He *et al.*, 2006b, 2007), strongly suggest a key role of Clp in the DSF regulatory pathway. The

speculation has been validated by the findings that deletion of *rpfF* or *rpfG* decreases the transcriptional expression of *clp* and overexpression of *clp* in the mutant $\Delta rpfF$ and $\Delta rpfG$, respectively, restores the production of EPS and extracellular enzymes to the wild-type level (He *et al.*, 2007). Moreover, microarray analysis shows that the DSF regulon and the Clp regulon are largely overlapping with similar expression patterns (He *et al.*, 2006b, 2007). Even though it is not clear how Clp detects and responds to the signal input from RpfG, the available lines of evidence have unequivocally established the important status of Clp in the DSF-signaling network (Fig. 3).

Clp mediates DSF signaling through a hierarchical regulatory network

Microarray analysis reveals that null mutation of Clp affects 299 genes at the transcriptional level (He *et al.*, 2007). The available data suggest that Clp modulates the transcriptional expression of this large set of genes by either direct activation or indirect regulation via other transcriptional regulators. Clp is homologous to the members of the Crp-FNR superfamily that are implicated in responding to a broad spectrum of intracellular and exogenous signals such as




Fig. 3. Schematic representation of the QS-signaling network in *Xcc*. The solid arrow indicates the demonstrated or predicted directed protein-protein interaction or directed signal modulation. The dashed arrows suggest a potential signal regulation pathway. As discussed in the text, it is likely that only a portion of the proteins with cyclic-di-GMP metabolic domains (GGDEF, EAL, and HD-GYP) may contribute to the change in the central pool of cyclic-di-GMP, with a subsequent effect on the transcriptional expression of the genes in the DSF regulon.

cAMP, anoxia, the redox state, oxidative and nitrosative stress, nitric oxide, carbon monoxide, 2-oxoglutarate, or temperature (Körner *et al.*, 2003). Within this family, the Crp of *E. coli* and the Vfr of *P. aeruginosa* have been well studied as global regulators affecting transcription of many genes directly or indirectly (Suh *et al.*, 2002; Martinez-Antonio & Collado-Vides 2003; Zheng *et al.*, 2004). Clp shares about 45% identity and 73% similarity to Crp (Dong & Ebright, 1992), which functions as a dimer in the form of a Crp–cAMP complex (Weber & Steitz, 1987; Lee *et al.*, 1994). The Clp dimer regulates transcriptional initiation by binding to a symmetrical DNA sequence (consensus sequence 5'-AAATGTGATCTAGATCACATTT-3'), located near or within the promoter regions (Zheng *et al.*, 2004). Similar to Crp and Vfr, Clp contains a conserved helix–turn–helix domain associated with DNA binding at the C-terminal and a putative cNMP-binding domain at the N-terminal (He *et al.*, 2007). It was shown that Clp has the same DNA-binding specificity as Crp at positions 5, 6, and 7 of the DNA half site (Dong & Ebright, 1992), and a consensus Clp-binding sequence (TGTGA-N6-TCACA) was thus suggested (Lee *et al.*, 2003). Using a computational algorithm with a confidence of 60% match to the nonvariable sequences of the putative Clp-binding motif, a total of 86 genes with at least one putative Clp-binding site at the promoter regions (arbitrarily defined as the 500 bp upstream of translation start codon ATG or GTG) were predicted out from the 299 Clp-regulated genes (He *et al.*, 2007). Most proteins or enzymes encoded by these 86 genes belong to the following functional groups: (1) extracellular enzymes, (2) EPS biosynthesis, (3) protein metabolism, and (4) membrane protein and transporter. These findings suggest that Clp may directly regulate the transcriptional expression of these genes (Fig. 3). Consistent with these findings, electrophoretic mobility shift assays and point mutation analysis confirm that Clp binds to the consensus Clp-binding sites at the promoter of *engXCA* gene, which encodes an extracellular cellulase (Hsiao *et al.*, 2005).

Further genetic and microarray analysis led to identification of a novel transcription factor FhrR, which shares < 20% amino acid identity with homologues in databases but contains a TetR family transcription factor domain, and a homologue of the Fur family transcription factor, designated as Zur for its function in the regulation of zinc uptake (Tang *et al.*, 2005; He *et al.*, 2007). The expression of both *zur* and *fhrR* is regulated by DSF and Clp and these two regulatory factors regulated different sets of genes within the Clp regulon (He *et al.*, 2007). These data suggest that Clp regulates the genes encoding flagella, Hrp, and ribosomal proteins through FhrR and the Clp-dependent regulation of iron uptake, multidrug resistance, and detoxification is mediated by Zur (Fig. 3). It is worth noting that DSF and Clp up- and downregulate more than a dozen genes

encoding various types of transcriptional factors (He *et al.*, 2006b, 2007). We could expect that further genetic and phenotypic analysis will reveal more regulatory elements within the DSF-Clp-regulatory networks.

Xcc biofilm formation is influenced by culture conditions and the DSF signal

One of the DSF-regulated biological functions is biofilm dispersal. In rich medium, mutation of the DSF synthase gene *rpfF*, or *rpfC/rpfG* encoding DSF sensing and signal transduction, results in formation of strain-dependent mild or extensive cell aggregates (biofilm) whereas the wild-type *Xcc* does not (Dow *et al.*, 2003; He *et al.*, 2006b). Consistently, exogenous addition of DSF at a physically relevant concentration to the *rpfF* deletion mutants grown in rich medium restores the wild-type planktonic growth mode (He *et al.*, 2006b). Interestingly, however, evidence is emerging that wild-type *Xcc* could also form a biofilm under favorable conditions. A recent study showed that in static cultures in minimum medium, wild-type *Xcc* form structured biofilms, whereas the *rpfF* and *rpfC* mutants (DSF overproducer and defective in DSF sensing) produce an unstructured biofilm (Torres *et al.*, 2007). Coinoculation of the wild-type strain 8004 with the *rpfC* mutant, which overproduces DSF, at a 1 : 1 ratio (but not at 4 : 1 ratio) abolishes the ability of 8004 to form a wild-type biofilm structure (Torres *et al.*, 2007). These findings suggest that *Xcc* may form different types of biofilms depending on the culture conditions and DSF signals.

In wild-type strain XC1, deletion of *rpfF* results in formation of cell aggregates comprising a few dozens to a few hundred cells, whereas deletion of *clp* alone does not affect DSF production, nor does it cause the formation of cell aggregates (He *et al.*, 2007). However, deletion of *clp* in the genetic background of the *rpfF* mutant leads to formation of much larger cell aggregates than the *rpfF* single deletion mutant, and exogenous addition of DSF to this double mutant disperses cell aggregations (He *et al.*, 2007). These phenotypes are puzzling, given that the transcriptional expression of *clp* is upregulated by DSF. Identification of other genes encoding for biofilm formation/dispersal and further analysis of the role of Clp in biofilm formation would facilitate characterization of these seemingly complicated regulatory mechanisms.

Biosynthesis of the DSF signal is autoregulated by a novel posttranslational regulatory mechanism

QS signals, such as AHL signals, are also widely known as autoinducers (Eberhard *et al.*, 1981; Fuqua & Greenberg, 2002), which are, in general, capable of autoregulating their own biosynthesis. In the QS system of *V. fischeri*, which is

considered as the paradigm of QS in Gram-negative bacteria, the AHL signal functions as a ligand to the transcription factor LuxR. At a lower cell density stage, each cell in the bacterial population produces a basal level of AHL signals via the AHL-synthase LuxI. At a sufficiently high population density, the accumulated AHL signals exceed a threshold level and interact with LuxR proteins; the resultant AHL-LuxR complex activates the transcriptional expression of QS-dependent genes as well as the *luxI* gene, leading to boosted production of AHL signals (for a review, see Dong *et al.*, 2007). This striking QS signal autoinduction mechanism enables bacteria to sense their population density, to synchronize the expression of the QS-regulon within the community in an effective way, and allows resetting of the whole QS circuit when a portion of bacterial cells are transferred to a new environment.

Evidence suggests that *Xcc* may have evolved a novel form of autoregulation mechanism that allows DSF to regulate its biosynthesis. Three *rpf* genes associated with DSF production have been identified in *Xcc*. Among them, *rpfF* and *rpfB* encode a putative enoyl CoA hydratase and a putative long-chain fatty acyl CoA ligase, respectively, and *rpfC* encodes a membrane-associated two-component sensor kinase (Barber *et al.*, 1997). Mutation of *rpfF* or *rpfB* abolishes DSF production and results in reduced virulence factor production (Barber *et al.*, 1997; He *et al.*, 2006b), whereas disruption of *rpfC* results in contrasting phenotypes. The *rpfC* mutant synthesizes about a 16-fold higher DSF signal than the wild-type *Xcc*, but produces significantly reduced virulence factors in a level similar to the *rpfF* mutant (Wang *et al.*, 2004). Among the two fatty acid metabolism enzymes, RpfF is particularly interesting as its expression in a non-DSF-producing bacterium, such as *E. coli*, confers DSF production activity (Y.-W. He & L.-H. Zhang, unpublished observation), suggesting that it is a key enzyme in the DSF biosynthesis pathway. However, while the time course analysis reveals that DSF production in *Xcc* increases proportionally following the increment of bacterial population density (Barber *et al.*, 1997; Wang *et al.*, 2004), the transcript level of *rpfF* remains more or less constant throughout growth and is not influenced by exogenous addition of DSF (Barber *et al.*, 1997; He *et al.*, 2006b). Moreover, even in the null mutant of *rpfC*, which produces abundant DSF signal, the level of *rpfF* transcripts shows only a moderate, less than twofold change (Slater *et al.*, 2000). These findings indicate that autoregulation of DSF biosynthesis unlikely occurs at the transcriptional level.

Two-component regulatory systems, consisting of a sensor and a cognate response regulator, are the predominant form of signal recognition and response coupling mechanism used by bacteria to sense and respond to various environmental cues, host signals, and bacterial QS signal molecules (Stock *et al.*, 2000; Wolanin *et al.*, 2002). The

orthodox sensor consists of only a sensing and an HK domain. Typically, recognition of a signal by the sensor triggers autophosphorylation at a histidine residue, followed by transfer of the phosphoryl group to an aspartate residue in the receiver domain of the cognate response regulator (Stock *et al.*, 2000), which leads to activation of transcriptional expression of the downstream genes within the regulon (Stock *et al.*, 2000). As discussed in the previous section, RpfC is a hybrid sensor consisting of multidomains including transmembrane, HK, receiver and HPT domains, and knocking out *rpfC* results in decreased virulence factor production but enhanced DSF biosynthesis. Several lines of evidence suggest that RpfC could modulate DSF production independent of its cognate response regulator RpfG and the conserved phosphorelay mechanism. First, unlike the *rpfC* mutant, the *rpfG* deletion mutant does not show a DSF overproduction phenotype and overexpression of RpfC in the *rpfGHC* triple-deletion mutant strongly represses DSF (Slater *et al.*, 2000). Second, overexpression of RpfG in the *rpfG/rpfC* double-deletion mutant restores the synthesis of virulence factors, but has no effect on repression of DSF biosynthesis (He, 2006). Third, site-directed mutagenesis of the three conserved amino acid residues involved in phosphorelay has no effect on DSF production (He *et al.*, 2006a). Furthermore, domain deletion analysis shows that the receiver domain of RpfC is essential for the RpfC-dependent repression of DSF biosynthesis. Overexpression of a short peptide of 107 aa within the receiver domain is sufficient to block the DSF production by the *rpfC* deletion mutant (He *et al.*, 2006a). These data collectively suggest a model in which uses two different mechanisms to regulate virulence factor and DSF production simultaneously, i.e., the hybrid sensor relies on the conserved phosphorelay mechanism and its cognate response regulator RpfG to induce the expression of virulence genes, and suppresses DSF biosynthesis by its receiver domain via an unknown mechanism.

Further analysis reveals that DSF may autoregulate its biosynthesis through a novel posttranslational mechanism involving RpfC–RpfF interaction. On the basis of the above findings, a construct expressing the receiver domain tagged with a 9-aa FLAG peptide was introduced into the *rpfC* deletion mutant. Coimmunoprecipitation and Far Western analysis demonstrate that REC binds tightly and specifically with RpfF, the key enzyme for DSF biosynthesis (He *et al.*, 2006a). These findings allow a rational speculation on how RpfC could induce the virulence factor synthesis by autoregulating DSF biosynthesis (Fig. 2). At a lower cell density or in an unconfined environment, the extracellular concentration of DSF is below a threshold and autophosphorylation of RpfC is not initiated. Unphosphorylated RpfC adopts a structural conformation that allows binding of RpfF to the REC domain, which limits the RpfF enzyme from interaction with the substrates provided by RpfB, thus

keeping the DSF synthesis at a basal level. Along with bacteria proliferation, the accumulated DSF signal interacts with and triggers autophosphorylation of RpfC, which results in a conformational change allowing release of RpfF, which leads to boosted DSF biosynthesis, and facilitating the four-step phosphorelay that activates RpfG and the downstream DSF regulon (He *et al.*, 2006a). In this manner, autoinduction of DSF biosynthesis is achievable without substantial elevation of *rpfF* transcription.

Interestingly, a recent study shows that in another *Xanthomonas* species, *Xanthomonas axonopodis* pv. *citri*, which contains conserved genes encoding RpfF, RpfC/RpfG, similar protein–protein interactions exist (Andrade *et al.*, 2006). The yeast two-hybrid and direct *in vitro* assays show that RpfC interacts with both RpfF and RpfG. The study also shows that the HD-GYP domain of RpfG interacts directly with several proteins containing the GGDEF domain. The finding suggests that this bacterial pathogen may also use the same posttranslational regulatory mechanism as *Xcc* for autoregulation of DSF production, and that RpfC and RpfG may regulate subsets of genes through protein–protein interactions. (Importantly, this need not involve alteration in gene expression.) Interestingly, a recent study shows that null mutation of RpfG results in decreased expression of three genes encoding the proteins containing GGDEF or EAL or both domains (Ryan *et al.*, 2007), although the mechanism of regulation remains unknown.

One core coupling to several sensors?

It becomes clear that the RpfC/RpfG two-component system couples the bacterial cell–cell communication to the bacterial intracellular regulatory networks through the global regulator Clp and the novel second messenger cyclic-di-GMP (Fig. 3). The intracellular level of cyclic-di-GMP is controlled through synthesis, catalyzed by the GGDEF protein domain, and degraded by the EAL or the HD-GYP domain (Tal *et al.*, 1998; D'Argenio *et al.*, 2004; Paul *et al.*, 2004; Christen *et al.*, 2005; Schmidt *et al.*, 2005; Ryan *et al.*, 2006). The finding that the HD-GYP domain of RpfG is a cyclic-di-GMP phosphodiesterase and that the functionality of RpfG is dependent on this enzyme activity (He, 2006; Ryan *et al.*, 2006) suggests that the intracellular level of cyclic-di-GMP is the key variable component that influences *Xcc* virulence and physiology. Intriguingly, besides RpfG, the genome of *Xcc* also encodes additional 36 proteins with the GGDEF, EAL, or HD-GYP domain (Ryan *et al.*, 2007). These proteins commonly contain other domains implicated in signaling sensing and receiving, such as receiver domain of a two-component response regulator (Stock *et al.*, 2000), GAF (Aravind & Ponting, 1997), HAMP (Aravind & Ponting, 1999), MHYT (Galperin *et al.*, 2001), and PAS domains (Taylor *et al.*, 2001; Wolanin *et al.*, 2002; Imamoto &

Kataoka, 2007). The contribution of each of these proteins to virulence, extracellular enzyme synthesis, and biofilm formation has been investigated using a panel of mutants (Ryan *et al.*, 2007). A total of 13 genes, including *rpfG*, are found with an effect on virulence on Chinese radish but only five of them significantly influence the production of extracellular enzyme virulence factors under *in vitro* conditions. It is similarly intriguing that only *rpfG* and *pdeA* seem to regulate biofilm formation and only *rpfG* controls motility under the conditions tested (Ryan *et al.*, 2007).

The above findings in *Xcc* are agreeable with the general patterns of cyclic-di-GMP signaling systems in other organisms (Jenal & Malone, 2006; Ryan *et al.*, 2007; Tamayo *et al.*, 2007), which have been summarized as follows: first, it is likely that not all the proteins with the GGDEF, EAL, or HD-GYP domains are involved in cyclic-di-GMP metabolism. Some of these proteins contain variations in the conserved GGDEF signature motifs (Ryan *et al.*, 2007). Evidence is accumulating that the proteins with noncanonical GGDEF domains have roles other than being directly associated with cyclic-di-GMP biosynthesis (Christen *et al.*, 2005; Kazmierczak *et al.*, 2006; Suzuki *et al.*, 2006). Second, specific signaling systems may modulate the intracellular levels of cyclic-di-GMP in a highly localized fashion, thus affecting the function of the proteins in the vicinity and allowing the pathogen to show distinct responses to different environmental cues. Alternatively, the difference in the relative enzyme activity and the expression level of the cyclic-di-GMP metabolic enzymes in various signaling systems could result in different changes in the level of cyclic-di-GMP. As the sensitivity of downstream regulatory systems to the change of cyclic-di-GMP (or related nucleotides) level may differ, these subtle differences would be translated to distinct bacterial responses to different environmental cues. Third, *Xcc* may be able to sense various cues of the environment and integrate the signal inputs by changing the cellular content of cyclic-di-GMP, and thus to modulate the bacterial physiology and virulence. A strong piece of supporting evidence is that several proteins with cyclic-di-GMP metabolic domains are implicated in the regulation of extracellular enzyme production in the presence of low oxygen tension and glutamine, respectively, which are the environmental conditions likely to be encountered by a vascular pathogen such as *Xcc* in the process of infection (Ryan *et al.*, 2007). For a clear understanding of the roles of these putative cyclic-di-GMP metabolic proteins in *Xcc* physiology and virulence, much work remains to be done.

Another key variable component in this virulence regulatory network is the global regulator Clp, whose intracellular level is influenced by DSF (He *et al.*, 2007). It is rather curious to note however, that, the *clp* transcripts, although decreased, still maintain a moderate level in the DSF null mutant Δ rpfF; this accompanies a moderate 28–40%

reduction in the production of EPS, protease, and cellulase, respectively. In contrast, the *clp* deletion mutant of XCC1 produces about 65–82% less EPS, protease, and cellulase, respectively, than its parent strain XCC1 (He *et al.*, 2007). Given that DSF may most likely influence the *clp* transcription via the enzyme activity of RpfG, as the deletion mutants Δ rpfF and Δ rpfG show a similar decrease in transcriptional expression of *clp* (He *et al.*, 2007), the factors that could change the intracellular level of cyclic-di-GMP may likely, but not necessarily, affect *clp* expression or influence its activity.

These findings suggest that Clp and cyclic-di-GMP may serve as the core of the regulatory network that modulates Xcc virulence, whose intracellular levels may change in response to different environmental cues, including those only found in host plants, and hence alter the expression patterns of downstream genes, including those coding for virulence factor production. In addition, as discussed above, RpfC and RpfG could also modulate subsets of genes through protein–protein interactions independent of Clp (Andrade *et al.*, 2006; He *et al.*, 2006a).

Intriguingly, the story does not seem to end here. A recent study shows that Xcc initiates expression of the *pip* gene encoding a proline iminopeptidase upon sensing an unknown signal(s) from the host plant, and mutation of *pip* results in a substantial decrease in bacterial virulence on host plant cabbage (Zhang *et al.*, 2007). Expression of *pip* requires a novel LuxR homologue designated as XccR that acts as a positive regulator. Mutation of *xccR* also resulted in significantly attenuated virulence of Xcc. Regardless of whether XccR/Pip modulates Xcc virulence through cyclic-di-GMP or Clp or other regulatory elements, this novel regulation mechanism clearly adds a new puzzle to the existing complexity of the Xcc virulence regulatory systems.

Conservation of the DSF-type QS system in other bacterial species

The availability of complete genomic sequences of over 300 bacterial species facilitates evaluation of the conservation of the DSF-signaling system among these bacteria. Using the key components of the DSF-signaling network (RpfF, RpfC, RpfG and Clp) as templates to BLAST the bacterial genome database in NCBI (<http://www.ncbi.nlm.nih.gov/genomes/lproks.cgi?view=1>), the counterparts of the four proteins showing high similarities (54–100% amino acid identity) are found from *X. campestris* pv. *vesicatoria*, *X. axonopodis* pv. *citri*, *Xanthomonas oryzae* pv. *oryzae*, *Xylella fastidiosa*, and *Stenotrophomonas maltophilia*. The homologues of RpfC, RpfG, and RpfF (34–51% amino acid identity) are also identified from the genomes of *Methylobacillus flagellatus*, *Thiobacillus denitrificans*, and *Leptospirillum* sp. The genes encoding the homologues of RpfF, RpfC, and RpfG

show a similar organization in their respective genomes; the only difference is the *rpfC* homologue UBAL79310316 of *Leptospirillum* sp. that shares the same orientation as UBAL79310317 (*rpfF*) (Fig. 4). The DSF synthase RpfF homologues with 28–37% amino acid identity are also found in the genomes of more than 60 species, for example, *Mariprofundus ferrooxydans* (ZP_01451794), *Sulfurovum* sp. (YP_001357875), *Burkholderia cenocepacia* (YP_837187), *Erythrobacter* sp. (ZP_01039845), *Yersinia mollaretii* (ZP_00825829), *Enterobacter sakazakii* (YP_001437677), *Serratia proteamaculans* (YP_001480964), and *Sphingomonas* sp. (ZP_01305181). However, no obvious homologues of the RpfC/RpfG two-component system could be identified from these genomes. These data suggest that the DSF-RpfC/RpfG-signaling system may very likely be conserved in *Xanthomonas*, *Xylella*, *Stenotrophomonas*, *Methylobacillus*, *Thiobacillus*, and *Leptospirillum*.

Consistent with the above bioinformatics information, DSF activity has been detected in a number of *Xanthomonas* spp., including *X. oryzae* pv. *oryzae* (Chatterjee & Sonti, 2002), *X. axonopodis* pv. *citri* (Jacques *et al.*, 2005), *X. fastidiosa* (Scarpari *et al.*, 2003; Newman *et al.*, 2004), *S. maltophilia* (Fouhy *et al.*, 2007), and *B. cenocepacia* (Boon *et al.*, 2008). The DSF extracts from *X. fastidiosa*, *S. maltophilia*, and *B. cenocepacia* restore the cellulase and EPS production in the *rpfF* mutant of Xcc, and vice versa, the DSF from Xcc restore the *rpfF*-dependent phenotypes in *S. maltophilia* and *X. fastidiosa* (Scarpari *et al.*, 2003; Fouhy *et al.*, 2007; Boon *et al.*, 2008), suggesting a conserved DSF structure among these bacterial pathogens. Recently, the DSF-like signaling molecule (BDSF) from *B. cenocepacia* has been purified and characterized as *cis*-dodecenoic acid (Boon *et al.*, 2008), which is highly similar to the DSF of Xcc but lacks only a methyl substitution. The structure of the DSF-like signal from *X. fastidiosa* has also been characterized tentatively as 12-methyl-tetradecanoic acid by MS analysis (Colnaghi Simionato *et al.*, 2007). Moreover, DSF (*cis*- Δ 2-11-methyl-dodecenoic acid) and several fatty acids or related structures have been identified from *S. maltophilia* (Huang & Wong, 2007a).

The general role of the DSF-signaling system in the modulation of virulence seems to be conserved in other bacterial species but the regulatory mechanisms and DSF-dependent traits may differ among taxa. In *X. oryzae* pv. *oryzae*, transposon insertional mutants of *rpfF* are deficient in virulence and exhibit an unusual tetracycline susceptibility phenotype and a growth deficiency under low iron conditions (Chatterjee & Sonti, 2002). The *rpfC* mutants of Xoo are defective in EPS production and attenuated in virulence on rice, but the mutation has no effect on extracellular enzyme biosynthesis (Tang *et al.*, 1996). As discussed in the previous section, *X. axonopodis* pv. *citri* seems to use similar mechanisms as Xcc to control DSF




Fig. 4. Conservation of the key genes involved in DSF synthesis, DSF perception, and DSF signal transduction in the genomes of different bacteria. RpfH is a protein showing significant peptide sequence similarity to the N-terminal part of RpfC. Its function in the DSF-signaling system remains unclear. Deletion of *rpfH* does not affect the virulence factor production. The numbers in the brackets indicate the percentages of identical amino acid compared with those in *Xcc* strain ATCC 33913. *Xcv*, *Xanthomonas campestris* pv. *vesicatoria*; *Xac*, *Xanthomonas axonopodis* pv. *citri*; *Xoo*, *Xanthomonas oryzae* pv. *oryzae*; *Xf*, *Xylella fastidiosa*; *Stm*, *Stenotrophomonas maltophilia*; *Mfla*, *Methylobacillus flagellatus*; *Tbd*, *Thiobacillus denitrificans*; and *Lep*, *Leptospirillum* sp Group II UBA. All sequences are retrieved from NCBI Microbial Genomes Resources (http://www.ncbi.nlm.nih.gov/genomes/MICROBES/microbial_taxtree.html).

production and DSF-dependent functions, because RpfC interacts with both RpfG and RpfF (Andrade *et al.*, 2006). In *X. fastidiosa*, DSF signaling has been found to influence the interactions of the pathogen with its insect vector and host plant. A more recent study shows that the DSF sensor RpfC is required for virulence and insect transmission of *X. fastidiosa* (Chatterjee *et al.*, 2008). Similar to *Xcc*, RpfC is involved in negative regulation of DSF biosynthesis by RpfF; *rpfC* mutants are deficient in virulence to grape. However, unlike in *Xcc*, mutation of *rpfF* increases virulence when the bacteria are mechanically inoculated into plants (Newman *et al.*, 2004). Reverse transcriptase (RT)-PCR analysis also reveals the contrasting effects of the *rpfC* and *rpfF* mutation on the expression pattern of a range of downstream genes including those encoding adhesins (Chatterjee *et al.*, 2008). Mutation of *rpfF* in *S. maltophilia* generates pleiotropic effects. The *rpfF* mutants show severely reduced motility, decreased production of extracellular protease, decreased tolerance to a range of antibiotics and heavy metals, and form cell aggregates (Fouhy *et al.*, 2007). In another report, deletion of *rpfF* or *rpfB* in *S. maltophilia* WR-C results in reduced transcription of *fecA*, which encodes a ferric citrate receptor that transports exogenous siderophore-ferric citrate from the environment into the bacterial periplasm, and addition of synthetic DSF restores the *FecA* expression (Huang & Wong, 2007b). Transposon inactivation of *crp*, which encodes a Clp homologue of *Xcc*, results in reduced *fecA* and *rpfF* transcriptional expression (Huang & Wong, 2007b).

Conclusions and future perspective

QS, in which bacterial cells communicate by means of small signal molecules, plays essential roles in the synchronization of gene expression and functional coordination among bacterial communities. The DSF signal, originally identified in *Xcc*, represents a novel class of the QS signal, that appears to be widely conserved in a range of bacterial species. In comparison with the classic AHL-type QS systems, the *Xcc* DSF-type QS-signaling system has evolved a unique auto-induction mechanism that modulates DSF biosynthesis, including a novel second messenger cyclic-di-GMP that facilitates coupling of QS signal sensing to the bacterial intracellular regulatory networks. Accumulating evidence also suggests that cyclic-di-GMP and the global regulator Clp, which contains a cNMP-binding domain, are the two key components that seem to serve as a core linking the responses to QS, environmental cues, and perhaps plant signals to the downstream genetic regulatory networks.

These exciting findings have laid down valuable frameworks for designing and developing novel strategies against bacterial infections, such as the quorum quenching or antipathogenic method against AHL-type QS systems (for reviews, see Williams, 2002; Zhang, 2003; Bjarnsholt & Givskov, 2007; Dong *et al.*, 2007). Similar to AHL-type signaling systems, DSF signaling is normally finely balanced during the disease process and disruption of such a fine balance could be a promising way for control of the disease.

Encouragingly, a recent study has identified a variety of bacteria isolates, belonging to the genera *Bacillus*, *Paenibacillus*, *Microbacterium*, *Staphylococcus*, and *Pseudomonas*, that show an obvious ability to inactivate the DSF signal molecules from *Xcc* (Newman *et al.*, 2008). Mutation of the *carAB* gene *Pseudomonas* spp. strain G, which encodes the biosynthesis of carbamoylphosphate, a precursor for pyrimidine and arginine biosynthesis, abolishes the DSF-inactivating ability of the bacterium (Newman *et al.*, 2008). Coinoculation of *Xcc* or *X. fastidiosa* with DSF-inactivating bacteria significantly reduces disease severity compared with plants inoculated with the pathogen alone (Newman *et al.*, 2008). Although the role of the DSF-inactivating enzyme in biocontrol remains to be further confirmed using corresponding mutants, this study has marked the beginning of developing quorum-quenching approaches against the DSF-mediated bacterial virulence. It can be sure that new strategies and agents will be identified and developed along with the progress in our understanding of DSF-signaling mechanisms in various pathogens.

In addition to the fascinating challenges of investigating the DSF-signaling mechanisms and networks in other bacterial pathogens, the current findings of DSF-signaling systems in *Xcc* have also presented many intriguing questions for further investigations in this model organism. For example, little is known about how the RpfC sensor may recognize the DSF signal; it is important to determine how RpfC may interact with RpfF and RpfG under various conditions; and it remains unclear how Clp may be influenced by cyclic-di-GMP and its metabolites, including pGpG and GMP. In addition, the finding that *Xcc* may grow either in a planktonic or in a biofilm form (Dow *et al.*, 2003; He *et al.*, 2006b), depending on the presence or absence of DSF QS signals, suggests that the pathogen could be an excellent model to study the mechanisms of transition of these two types of common bacterial growth forms. Moreover, it remains a fascinating challenge to determine how the various cyclic-di-GMP metabolic enzymes are activated and hence influence the bacterial virulence and physiology. Furthermore, the fascinating finding that DSF-like signals could influence the biological functions of eukaryotic fungal pathogens at biologically relevant concentrations (Wang *et al.*, 2004; Boon *et al.*, 2008) may substantially broaden the scope of investigation on DSF-type QS systems.

References

- Andrade MO, Alegria MC, Guzzo CR, Docena C, Rosa MC, Ramos CH & Farah CS (2006) The HD-GYP domain of RpfG mediates a direct linkage between the Rpf quorum-sensing pathway and a subset of diguanylate cyclase proteins in the phytopathogen *Xanthomonas axonopodis* pv. *citri*. *Mol Microbiol* **62**: 537–551.
- Anon (1978) *Distribution Maps of Plant Diseases No. 136*. 4th edn. Commonwealth Mycological Institute, Kew, Surrey, UK.
- Aravind L & Koonin EV (1998) The HD domain defines a new superfamily of metal-dependent phosphohydrolases. *Trends Biochem Sci* **23**: 469–472.
- Aravind L & Ponting CP (1997) The GAF domain: an evolutionary link between diverse phototransducing proteins. *Trends Biochem Sci* **22**: 458–459.
- Aravind L & Ponting CP (1999) The cytoplasmic helical linker domain of receptor histidine kinase and methyl-accepting proteins is common to many prokaryotic signalling proteins. *FEMS Microbiol Lett* **176**: 111–116.
- Barber CE, Tang JL, Feng JX, Pan MQ, Wilson TJG, Slater H, Dow JM, Williams P & Daniels MJ (1997) A novel regulatory system required for pathogenicity of *Xanthomonas campestris* is mediated by a small diffusible signal molecule. *Mol Microbiol* **24**: 556–566.
- Bassler BL & Losick R (2006) Bacterially speaking. *Cell* **125**: 237–246.
- Bjarnsholt T & Givskov M (2007) Quorum-sensing blockade as a strategy for enhancing host defences against bacterial pathogens. *Phil Trans R Soc B* **362**: 1213–1222.
- Blevess S, Soscia C, Nogueira-Orlandi P, Lazdunski A & Filloux A (2005) Quorum sensing negatively controls type III secretion regulon expression in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* **187**: 3898–3902.
- Boon C, Deng Y, Wang LH, He YW, Xu JL, Fan Y, Pan SQ & Zhang LH (2008) A novel DSF-like signal from *Burkholderia cenocepacia* interferes with *Candida albicans* morphological transition. *ISME J* **2**: 27–36.
- Chang WH, Lee MC, Yang MT & Tseng YH (2005) Expression of heat-shock genes *groESL* in *Xanthomonas campestris* is upregulated by CLP in an indirect manner. *FEMS Microbiol Lett* **243**: 365–372.
- Chatterjee S & Sonti RV (2002) *rpfF* mutants of *Xanthomonas oryzae* pv. *oryzae* are deficient for virulence and growth under low iron conditions. *Mol Plant-Microbe Interact* **15**: 463–471.
- Chatterjee S, Wistrom C & Lindow SE (2008) A cell-cell signaling sensor is required for virulence and insect transmission of *Xylella fastidiosa*. *Proc Natl Acad Sci USA* **105**: 2670–2675.
- Chen X, Schauder S, Potier N, Van Dorselaer A, Pelczar I, Bassler BL & Hughson FM (2002) Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* **415**: 545–549.
- Christen M, Christen B, Folcher M, Schauerte A & Jenal U (2005) Identification and characterization of a cyclic di-GMP-specific phosphodiesterase and its allosteric control by GTP. *J Biol Chem* **280**: 30829–30837.
- Chun W, Cui J & Poplawsky A (1997) Purification, characterization and biological role of a pheromone produced by *Xanthomonas campestris* pv. *campestris*. *Physiol Mol Plant Pathol* **51**: 1–14.
- Colnaghi Simionato AV, da Silva DS, Lambais MR & Carrilho E (2007) Characterization of a putative *Xylella fastidiosa*

- diffusible signal factor by HRGC-EI-MS. *J Mass Spectrom* **42**: 490–496.
- D'Argenio DA & Miller SI (2004) Cyclic-di-GMP as a bacterial second messenger. *Microbiology* **150**: 2497–2502.
- de Crecy-Lagard V, Glaser P, Lejeune P, Sismeiro O, Barber CE, Daniels MJ & Danchin A (1990) A *Xanthomonas campestris* pv. *campestris* protein similar to catabolite activation factor is involved in regulation of phytopathogenicity. *J Bacteriol* **172**: 5877–5883.
- da Silva AC, Ferro JA, Reinach FC *et al.* (2002) Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* **417**: 459–463.
- Dong Q & Ebricht RH (1992) DNA binding specificity and sequence of *Xanthomonas campestris* catabolite gene activator protein-like protein. *J Bacteriol* **174**: 5457–5461.
- Dong YH, Wang LH & Zhang LH (2007) Quorum quenching microbial infections – mechanisms and implications. *Phil Trans R Soc B* **362**: 1201–1211.
- Dow JM & Daniels MJ (1994) Pathogenicity determinants and global regulation of pathogenicity of *Xanthomonas campestris* pv. *campestris*. *Curr Top Microbiol Immunol* **192**: 29–41.
- Dow JM, Crossman L, Findlay K, He YQ, Feng JX & Tang JL (2003) Biofilm dispersal in *Xanthomonas campestris* is controlled by cell–cell signaling and is required for full virulence to plants. *Proc Natl Acad Sci USA* **100**: 10995–11000.
- Eberhard A, Burlingame AL, Eberhard C, Kenyon GL, Nealson KH & Oppenheimer NJ (1981) Structural identification of autoinducer of *Photobacterium fischeri* luciferase. *Biochem* **20**: 2444–2449.
- Flavier AB, Clough SJ, Schell MA & Denny TP (1997) Identification of 3-hydroxypalmitic acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*. *Mol Microbiol* **26**: 251–259.
- Fouhy Y, Scanlon K, Schouest K, Spillane C, Crossman L, Avison MB, Ryan RP & Dow JM (2007) Diffusible signal factor-dependent cell–cell signaling and virulence in the *Nosocomial* pathogen *Stenotrophomonas maltophilia*. *J Bacteriol* **189**: 4964–4968.
- Fuqua C & Greenberg EP (2002) Listening in on bacteria: acyl-homoserine lactone signalling. *Nat Rev Mol Cell Biol* **3**: 685–695.
- Galperin MY, Natale DA, Aravind L & Koonin EV (1999) A specialized version of the HD hydrolase domain implicated in signal transduction. *J Mol Microbiol Biotechnol* **1**: 303–305.
- Galperin MY, Gaidenko TA, Mulikidjanian AY, Nakano M & Price CW (2001) MHYT, a new integral membrane sensor domain. *FEMS Microbiol Lett* **205**: 17–23.
- García-Ochoa F, Santos VE, Casas JA & Gómez E (2000) Xanthan gum: production, recovery, and properties. *Biotechnol Adv* **18**: 549–579.
- Ghosh P (2004) Process of protein transport by the type III secretion system. *Microbiol Mol Biol Rev* **68**: 771–795.
- Hayward AC (1993) The hosts of *Xanthomonas*. *Xanthomonas* (Swings G & Civerolo EL, eds), pp. 1–119. Chapman and Hall, London.
- He YW (2006) Cell–cell communication in *Xanthomonas campestris* pv. *campestris*. PhD Thesis, Library of National University of Singapore (<http://libpweb1.nus.edu.sg/lion//wtheses.html>), Singapore.
- He YW, Wang C, Zhou L, Song H, Dow JM & Zhang LH (2006a) Dual signaling functions of the hybrid sensor kinase RpfC of *Xanthomonas campestris* involve either phosphorelay or receiver domain–protein interaction. *J Biol Chem* **281**: 33414–33421.
- He YW, Xu M, Lin K *et al.* (2006b) Genome scale analysis of diffusible signal factor regulon in *Xanthomonas campestris* pv. *campestris*: identification of novel cell–cell communication-dependent genes and functions. *Mol Microbiol* **59**: 610–622.
- He YW, Ng AY, Xu M, Lin K, Wang LH, Dong YH & Zhang LH (2007) *Xanthomonas campestris* cell–cell communication involves a putative nucleotide receptor protein Clp and a hierarchical signalling network. *Mol Microbiol* **64**: 281–292.
- Hsiao YM & Tseng YH (2002) Transcription of *Xanthomonas campestris prt1* gene encoding protease 1 increases during stationary phase and requires global transcription factor Clp. *Biochem Biophys Res Commun* **295**: 43–49.
- Hsiao YM, Liao HY, Lee MC, Yang TC & Tseng YH (2005) Clp upregulates transcription of *engA* gene encoding a virulence factor in *Xanthomonas campestris* by direct binding to the upstream tandem Clp sites. *FEBS Lett* **579**: 3525–3533.
- Huang TP & Wong AC (2007a) Extracellular fatty acids facilitate flagella-independent translocation by *Stenotrophomonas maltophilia*. *Res Microbiol* **158**: 702–711.
- Huang TP & Wong AC (2007b) A cAMP receptor protein regulated cell–cell communication system mediates expression of a FecA homologue in *Stenotrophomonas maltophilia*. *Appl Environ Microbiol* **73**: 5034–5040.
- Hueck CJ (1998) Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol Mol Biol Rev* **62**: 379–433.
- Hugouvieux V, Barber CE & Daniels MJ (1998) Entry of *Xanthomonas campestris* pv. *campestris* into hydathodes of *Arabidopsis thaliana* leaves: a system for studying early infection events in bacterial pathogenesis. *Mol Plant–Microbe Interact* **11**: 537–543.
- Imamoto Y & Kataoka M (2007) Structure and photoreaction of photoactive yellow protein, a structural prototype of the PAS domain superfamily. *Photochem Photobiol* **83**: 40–49.
- Jacques MA, Josi K, Darrasse A & Samson R (2005) *Xanthomonas axonopodis* pv. *phaseoli* var. *fuscans* is aggregated in stable biofilm population sizes in the phyllosphere of field-grown beans. *Appl Environ Microbiol* **71**: 2008–2015.
- Jenal U & Malone J (2006) Mechanisms of cyclic-di-GMP signaling in bacteria. *Annu Rev Genet* **40**: 385–407.
- Kazmierczak BI, Lebron MB & Murray TS (2006) Analysis of FimX, a phosphodiesterase that governs twitching motility in *Pseudomonas aeruginosa*. *Mol Microbiol* **60**: 1026–1043.
- Kennedy JF & Bradshaw IJ (1984) Production, properties and applications of xanthan. *Prog Ind Microbiol* **19**: 319–371.

- Körner H, Sofia HJ & Zumft WG (2003) Phylogeny of the bacterial superfamily of Crp-Fnr transcription regulators: exploiting the metabolic spectrum by controlling alternative gene programs. *FEMS Microbiol Rev* **27**: 559–592.
- Lee EJ, Glasgow J, Leu SF, Belduz AO & Harman JG (1994) Mutagenesis of the cyclic AMP receptor protein of *Escherichia coli*: targeting position 83, 127 and 128 of the cyclic nucleotide binding pocket. *Nucleic Acids Res* **22**: 2894–2901.
- Lee MC, Weng SF & Tseng YH (2003) Flagellin gene *fliC* of *Xanthomonas campestris* is upregulated by transcription factor Clp. *Biochem Biophys Res Commun* **307**: 647–652.
- Leela JK & Sharma G (2000) Studies on xanthan production from *Xanthomonas campestris*. *Bioproc Biosystems Eng* **23**: 687–689.
- Leyns F, De Cleene M, Swings J & De Ley J (1984) The host range of the genus *Xanthomonas*. *Bot Rev* **50**: 308–355.
- Lilly VG, Watson HA & Leach JG (1958) Bacterial polysaccharide, II. Laboratory-scale production of polysaccharides by species of *Xanthomonas*. *Appl Microbiol* **6**: 105–108.
- Lopes CA & Quezado-Soares AM (1997) *Doenças bacterianas das hortaliças: diagnose e controle*. Embrapa, Serviço de Produção de Informação, Brasília.
- Martinez-Antonio A & Collado-Vides J (2003) Identifying global regulators in transcriptional regulatory networks in bacteria. *Curr Opin Microbiol* **6**: 482–489.
- Miller ST, Xavier KB, Campagna SR, Taga ME, Semmelhack MF, Bassler BL & Hughson FM (2004) *Salmonella typhimurium* recognizes a chemically distinct form of the bacterial quorum-sensing signal AI-2. *Mol Cell* **15**: 677–687.
- Morris ER, Rees DA, Walkinsaw MD & Darke A (1977) Order–disorder transition for a bacterial polysaccharide in solution. *J Mol Biol* **110**: 1–16.
- Newman KL, Almeida RPP, Purcell AH & Lindow SE (2004) Cell–cell signaling controls *Xylella fastidiosa* interactions with both insects and plants. *Proc Natl Acad Sci USA* **101**: 1737–1742.
- Newman KL, Chatterjee S, Ho KA & Lindow SE (2008) Virulence of plant pathogenic bacteria attenuated by degradation of fatty acid cell-to-cell signaling factors. *Mol Plant–Microbe Interact* **21**: 326–334.
- Onsando JM (1992) Black rot of crucifers. *Diseases of Vegetables and Oil Seed Crops, Vol. II* (Chaube HS, Kumar J, Mukhopadhyay AN & Singh US, eds), pp. 243–252. Prentice Hall, New Jersey.
- Paul R, Weiser S, Amiot NC, Chan C, Schirmer T, Giese B & Jenal U (2004) Cell cycle-dependent dynamic localization of a bacterial response regulator with a novel di-guanylate cyclase output domain. *Genes Dev* **18**: 715–727.
- Poplawsky AR & Chun W (1997) *pigB* determines a diffusible factor needed for extracellular polysaccharide slime and xanthomonadin production in *Xanthomonas campestris* pv. *campestris*. *J Bacteriol* **179**: 439–444.
- Poplawsky AR & Chun W (1998) *Xanthomonas campestris* pv. *campestris* requires a functional *pigB* for epiphytic survival and host infection. *Mol Plant–Microbe Interact* **11**: 466–475.
- Poplawsky AR, Chun W, Slater H, Daniels MJ & Dow JM (1998) Synthesis of extracellular polysaccharide, extracellular enzymes, and xanthomonadin in *Xanthomonas campestris*: evidence for the involvement of two intercellular regulatory signals. *Mol Plant–Microbe Interact* **11**: 68–70.
- Poplawsky AR, Urban SC & Chun W (2000) Biological role of xanthomonadin pigments in *Xanthomonas campestris* pv. *campestris*. *Appl Environ Microbiol* **66**: 5123–5127.
- Poplawsky AR, Walters DM, Rouviere PE & Chun W (2005) A gene for a dioxygenase-like protein determines the production of the DF signal in *Xanthomonas campestris* pv. *campestris*. *Mol Plant Pathol* **6**: 653–657.
- Qian W, Jia Y, Ren SX *et al.* (2005) Comparative and functional genomic analyses of the pathogenicity of phytopathogen *Xanthomonas campestris* pv. *campestris*. *Genome Res* **15**: 757–767.
- Römmling U, Gomelsky M & Galperin MY (2005) C-di-GMP: the dawning of a novel bacterial signaling system. *Mol Microbiol* **57**: 629–639.
- Ryan RP, Fouhy Y, Lucey JF *et al.* (2006) Cell–cell signaling in *Xanthomonas campestris* involves an HD-GYP domain protein that functions in cyclic di-GMP turnover. *Proc Natl Acad Sci USA* **103**: 6712–6717.
- Ryan RP, Fouhy Y, Lucey JF, Jiang BL, He YQ, Feng JX, Tang JL & Dow JM (2007) Cyclic di-GMP signalling in the virulence and environmental adaptation of *Xanthomonas campestris*. *Mol Microbiol* **63**: 429–442.
- Scarpari LM, Lambais MR, Silva DS, Carraro DM & Carrer H (2003) Expression of putative pathogenicity-related genes in *Xylella fastidiosa* grown at low and high cell density conditions *in vitro*. *FEMS Microbiol Lett* **222**: 83–92.
- Schmidt AJ, Ryjenkov DA & Gomelsky M (2005) Ubiquitous protein domain EAL encodes cyclic diguanylate-specific phosphodiesterase: enzymatically active and inactive EAL domains. *J Bacteriol* **187**: 4774–4781.
- Schuster M, Lostroh CP, Ogi T & Greenberg EP (2003) Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. *J Bacteriol* **185**: 2066–2079.
- Slater H, Alvarez-Morales A, Barber CE, Daniels MJ & Dow JM (2000) A two-component system involving an HD-GYP domain protein links cell–cell signaling to pathogenicity gene expression in *Xanthomonas campestris*. *Mol Microbiol* **38**: 986–1003.
- Stock AM, Robinson VL & Goudreau PN (2000) Two-component signal transduction. *Ann Rev of Biochem* **69**: 183–215.
- Suh SJ, Runyen-Janecky LJ, Maleniak TC, Hager P, MacGregor CH, Zielinski-Mozny NA, Phibbs PV Jr & West SE (2002) Effect of *vfr* mutation on global gene expression and catabolite repression control of *Pseudomonas aeruginosa*. *Microbiology* **148**: 1561–1569.
- Sutherland IW (1998) Novel and established applications of microbial polysaccharides. *Tib Tech* **16**: 41–46.
- Suzuki K, Babitzke P, Kushner SR & Romeo T (2006) Identification of a novel regulatory protein (CsrD) that targets

- the global regulatory RNAs CsrB and CsrC for degradation by RNase E. *Genes Dev* **20**: 2605–2617.
- Tal R, Wong HC, Calhoon R *et al.* (1998) Three *cdg* operons control cellular turnover of cyclic-di-GMP in *Acetobacter xylinum*: genetic organization and occurrence of conserved domains in isoconymes. *J Bacteriol* **180**: 4416–4425.
- Tamayo R, Pratt JT & Camilli A (2007) Roles of cyclic diguanylate in the regulation of bacterial pathogenesis. *Annu Rev Microbiol* **61**: 131–148.
- Tang JL, Liu YN, Barber CE, Dow JM, Wootton JC & Daniels MJ (1991) Genetic and molecular analysis of a cluster of *rpf* genes involved in positive regulation of synthesis of extracellular enzymes and polysaccharide in *Xanthomonas campestris* pv. *campestris*. *Mol Gen Genet* **226**: 409–417.
- Tang JL, Feng JX, Li QQ, Wen HX, Zhou DL, Wilson TJ, Dow JM, Ma QS & Daniels MJ (1996) Cloning and characterization of the *rpfC* gene of *Xanthomonas oryzae* pv. *oryzae*: involvement in exopolysaccharide production and virulence to rice. *Mol Plant–Microbe Interact* **9**: 664–666.
- Tang DJ, He YQ, Feng JX, He BR, Jiang BL, Lu GT, Chen B & Tang JL (2005) *Xanthomonas campestris* pv. *campestris* possesses a single gluconeogenic pathway that is required for virulence. *J Bacteriol* **187**: 6231–6237.
- Taylor BL, Rebbapragada A & Johnson MS (2001) The PAD-PAS domain as a sensor for behavioral responses in *Escherichia coli*. *Antioxid Redox Signaling* **3**: 867–879.
- Torres PS, Malamud F, Rigano LA, Russo DM, Marano MR, Castagnaro AP, Zorreguieta A, Bouarab K, Dow JM & Vojnov AA (2007) Controlled synthesis of the DSF cell–cell signal is required for biofilm formation and virulence in *Xanthomonas campestris*. *Environ Microbiol* **9**: 2101–2109.
- Vendeville A, Winzer K, Heurlier K, Tang CM & Hardie KR (2005) Making ‘sense’ of metabolism: autoinducer-2, LuxS and pathogenic bacteria. *Nat Rev Microbiol* **3**: 383–396.
- Wang LH, He YW, Gao YF *et al.* (2004) A bacterial cell–cell communication signal with cross-kingdom structural analogues. *Mol Microbiol* **51**: 903–912.
- Weber IT & Steitz TA (1987) Structure of a complex of catabolite gene activator protein and cyclic AMP refined at 2.5 resolution. *J Mol Biol* **198**: 311–326.
- Whitehead NA, Barnard AM, Slater H, Simpson NJ & Salmond GP (2001) Quorum-sensing in Gram-negative bacteria. *FEMS Microbiol Rev* **25**: 365–404.
- Williams PH (1980) Black rot: a continuing threat to world crucifers. *Plant Dis* **64**: 736–742.
- Williams P (2002) Quorum sensing: an emerging target for antibacterial chemotherapy? *Expert Opin Ther Targets* **6**: 1–18.
- Williams P, Winzer K, Chan WC & Cámara M (2007) Look who’s talking: communication and quorum sensing in the bacterial world. *Philos Trans R Soc Lond B Biol Sci* **362**: 1119–1134.
- Wolanin PM, Thomason PA & Stock JB (2002) Histidine protein kinases: key signal transducers outside the animal kingdom. *Genome Biol* **3**: 1–8.
- Xavier KB & Bassler BL (2003) LuxS quorum sensing: more than just a numbers game. *Curr Opin Microbiol* **6**: 191–197.
- Yun MH, Torres PS, Oirdi ME, Rigano LA, Gonzalez-Lamothe R, Marano MR, Castagnaro AP, Dankert MA, Bouarab K & Vojnov AA (2006) Xanthan induces plant susceptibility by suppressing callose deposition. *Plant Physiol* **141**: 178–187.
- Zhang LH (2003) Quorum quenching and proactive host defense. *Trends Plant Sci* **8**: 238–244.
- Zhang LH & Dong YH (2004) Quorum sensing and signal interference: diverse implications. *Mol Microbiol* **53**: 1563–1571.
- Zhang L, Jia Y, Wang L & Fang R (2007) A proline iminopeptidase gene upregulated *in planta* by a LuxR homologue is essential for pathogenicity of *Xanthomonas campestris* pv. *campestris*. *Mol Microbiol* **65**: 121–136.
- Zheng D, Constantinidou C, Hobman JL & Minchin SD (2004) Identification of the CRP regulon using *in vitro* and *in vivo* transcriptional profiling. *Nucleic Acids Res* **32**: 5874–5893.