

IMMEDIATE ONLINE ACCEPTED (IOA)
ARTICLE

This article presented here has been peer reviewed and accepted for publication in *CCS Chemistry*. The present version of this manuscript has been posted at the request of the author prior to copyediting and composition and will be replaced by the final published version once it is completed. The DOI will remain unchanged.

IOA Posting Date: December 10 2024

TITLE: Discovery of a novel family of Gram-negative-active cationic cyclolipopeptides by motif search-guided chemical synthesis

AUTHORS: Fan Zhang, Fei Zhang, Xuchang Yu, Runze Sun, Yuzhu Wu, Yiwen Zhou and Lei Li

DOI: 10.31635/ccschem.024.202404925

CITE THIS: *CCS Chem.* 2024, Just Accepted. DOI: 10.31635/ccschem.024.202404925

1
2
3
4
5
6 **Discovery of a novel family of Gram-negative-active cationic cyclolipopeptides by motif**
7
8 **search-guided chemical synthesis**
9

10 Fan Zhang^{1#}, Fei Zhang^{1#}, Xuchang Yu^{1,2#}, Runze Sun¹, Yuzhu Wu¹, Yiwen Zhou¹ and Lei Li^{1*}
11
12

13 ¹ State Key Laboratory of Microbial Metabolism and School of Life Sciences and Biotechnology, Shanghai Jiao
14 Tong University, 800 Dongchuan Rd., Shanghai, 200240
15

16 ² State Key Laboratory of Bioreactor Engineering and School of Biotechnology, East China University of Science
17 and Technology, 130 Meilong Rd., Shanghai 200237
18

19 *Corresponding Author: Lei Li: lei.li@sjtu.edu.cn
20
21
22

23
24 **Abstract**
25

26
27 Emerging antimicrobial resistance is a global public health crisis, which necessitates the development of
28 antibiotics with novel structural scaffolds and modes-of-action (MOAs). Cationic cyclolipopeptides have been
29 increasingly recognized as a kind of promising therapeutics against Gram-negative pathogens. By two-round
30 motif searching of 8,982 sequenced genomes from 42 underexplored Gram-negative bacterial genera, here we
31 identified a novel family of cationic cyclolipopeptide biosynthetic gene clusters (BGCs). Then, a synthetic-
32 bioinformatic natural product (syn-BNP) approach that relies on structure prediction with high accuracy
33 followed by chemical synthesis, was used to rapidly access their predicted products with six positively charged
34 amino acids and nine D-amino acids out of 12 building blocks, thus generating chisopeptin and its five
35 congeners. All of them showed good activities against a panel of Gram-negative pathogens with MIC of 2-8
36 µg/mL. Furthermore, chisopeptin was found to have a unique MOA to kill *Klebsiella pneumoniae* by possibly
37 targeting the inner membrane protein YejM that regulates lipopolysaccharide biogenesis. The study of
38 chisopeptin-type structures inspired by underexplored BGC families provides a productive approach to develop
39 novel cyclolipopeptide-based therapeutics.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

Keywords

Anti-microbial resistance, underexplored sources, natural products, cationic cyclolipopeptides, chemical synthesis, YeJM

Introduction

The emergence of multidrug-resistant Gram-negative pathogens has become a major threat to global public health and necessitates the development of novel antibiotics with diverse modes of action (MOAs).¹⁻³ The chemical and mechanistic diversity of bacterial non-ribosomal peptides (NRPs) encoded by giant NRP synthetases (NRPSs) have been known as promising natural products for novel antibiotic discovery.⁴⁻⁷ Multiple chemical modifications, such as cyclization, introduction of positively charged amino acids (PCAAs) or D-amino acids (D-AAs) and *N*-terminal fatty acid acylation, are usually used to optimize the physicochemical and proteolytic stability profiles of peptides, further improving their therapeutic potentials.⁸ Particularly, cationic lipopeptides (CLPs) with at least two PCAAs, showed unique chemical properties for combating Gram-negative pathogens and have been developed for a series of clinically used antibiotics (i.e., colistin and polymyxin B).⁹⁻

11

In the last ten years, underexplored Gram-negative bacteria have been increasingly recognized as rich sources of NRPs for the discovery of novel CLP-type antibiotics.¹²⁻¹⁵ In our previous study, 8982 Gram-negative bacterial genomes from 42 underexplored genera were screened for novel CLP biosynthetic gene clusters (BGCs).¹⁶ A linear CLP chospeptin was rapidly identified by structure prediction followed by solid-phase peptide synthesis, which showed good activities against different colistin-resistant Gram-negative pathogens by directly targeting lipopolysaccharide (LPS).¹⁶ However, although the eight novel CLP BGCs were selected but we just synthesized their inspired linear products, possibly explaining why only chospeptin showed good antibacterial activity (Figure 1 and Table S1).¹⁶ We proposed that the other seven CLP BGCs might encode cyclolipopeptides considering some hydrophilic AAs (i.e., Dap, Dab, Ser and Thr) as nucleophiles could realize peptide cyclization

through the C-terminal carboxyl. Actually, cyclization has emerged as the most widely used strategy to optimize the bioactivity of natural or synthetic peptides.¹⁷ Considering more than 40 potential cyclolipopeptides inspired by the other seven CLP BGCs need to be synthesized, another AA modification (introduction of D-AAs) was used for the second-round motif search (Table S1). Natural peptides usually comprise L-amino acids (L-AAs), which are vulnerable to enzyme degradation. By contrast, introduction of D-AAs could often optimize the proteolytic stability of peptides, thus improving their bioactivity.^{8,18,19} While multiple D-AAs in a single peptide was used for the second-round search, we found an interesting CLP2 BGC from *Chitinophaga solisilvae* CGMCC 1.12462, whose predicted p




Figure 1 | Two-round motif search for novel cationic lipopeptides (CLPs) with multiple D-amino acids from underexplored Gram-negative bacteria. One interesting CLP BGC whose encoded peptide contains nine D-amino acids was selected for structure prediction followed by solid-phase peptide synthesis as either linear or cyclic structures.

Experimental Methods

Chemical reagents, consumables and instruments

Standard *N*-Fmoc amino acids were purchased from GL Biochem (Shanghai, CN). Lipopolysaccharides (LPS) from *Escherichia coli* O55:B5 was purchased from Beijing InnoChem (Beijing, CN). GlcNAC, lipoteichoic acid,

1
2
3
4 muramyl dipeptide, MurNAC, pentapeptide and peptidoglycan were purchased from Bidepharm (Shanghai,
5
6 CN). Reagents for solid-phase peptide synthesis (SPPS), including 2-chlorotrityl resin, dichloromethane (DCM),
7
8 *N,N*-diisopropylethylamine (DIPEA), 4-dimethylaminopyridine (DMAP), *N,N*-dimethylformamide (DMF), *O*-(7-
9
10 azabenzotriazol-1-yl)-*N,N,N'*-tetramethyluronium hexafluorophosphate (HATU), hexafluoroisopropanol
11
12 (HFIP), (7-azabenzotriazol-1-yloxy) tripyrrolidinophosphonium hexafluorophosphate (PyAOP) and
13
14 trifluoroacetic acid (TFA), were purchased from GL Biochem (Shanghai, CN) or Shanghai Titan (Shanghai, CN).
15
16

17
18 An Agilent 1260 Series HPLC with UV detection and equipped with an XBridge Prep C18 130 Å column (10 x 150
19
20 mm, 5 µm) was used for peptide purification. For all liquid chromatography, solvent A = H₂O and solvent B =
21
22 CH₃CN. An Agilent 1290 Series HPLC coupled to a 6546 Series QTOF mass spectrometer, equipped with a
23
24 Phenomenex Kinetex PS C18 100 Å column (2.1 x 50 mm, 2.6 µm) and controlled by Masshunter software, was
25
26 used to acquire the HPLC-HRMS data of synthetic peptides. A Bruker Avance DMX 700 MHz spectrometer
27
28 equipped with cryogenic probes was used to acquire the ¹H, ¹³C, DEPT135, COSY, HMBC and HSQC NMR spectra
29
30 of synthetic peptides. All spectra were recorded at 25 °C in DMSO-*d*₆. Chemical shift values are reported in
31
32 parts in million (ppm) and referenced to residual solvent signals: 2.50 ppm (¹H) and 39.52 ppm (¹³C).
33
34
35

36 **Solid Phase Peptide Synthesis (SPPS)**

37
38
39 Standard Fmoc chemistry-based solid-phase peptide synthesis methods were used to synthesize peptides on
40
41 2-chlorotrityl chloride resin. Peptide synthesis started from the last AA of each peptide, which was loaded on
42
43 2-chlorotrityl resin (0.3 g, 0.552 mmol/g) and was swollen in DCM for 20 min, drained as well as washed with
44
45 DMF (3 mL, 3x). By using Fmoc-protected amino acids (2 equiv., relative to resin loading) mixed with HATU (2
46
47 equiv.) and DIPEA (2 equiv.) in DMF (5 mL), individual amino acids were coupled step by step. Finally, coupling
48
49 reactions were performed for 1 hr with occasional swirling and washed with DMF (3 mL, 3x).
50
51
52
53
54
55
56
57
58

1
2
3
4
5 For the synthesis of synCLP2-L, Fmoc-deprotection was done using 20% piperidine in DMF (3 mL) for 7 min and
6
7 repeated twice. The resin was washed with DMF (3 mL, 5x) and then coupled with a subsequent *N*-Fmoc amino
8
9 acid. Then, tetradecanoic acid was coupled into the end of linear peptide.

10
11
12 For the synthesis of synCLP2-C1, synCLP2-C7 (chisopeptin) and chisopeptin congeners, the alloc group of Dap
13
14 was firstly removed by two rounds of palladium [Pd(PPh₃)₄]-catalyzed deprotection in DCM. In each round of
15
16 deprotection, 0.25 equivalent palladium and 12 equivalent phenylsilane were added to resins suspended in DCM
17
18 (3 mL) and reacted for 1 hr. Then, 2-chlorotriptyl resin was removed by 20% HFIP in DCM for 2 hrs. Soluble
19
20 peptides were further cyclized using PYAOP (8 equivalent) and DIPEA (30 equivalent) in DMF and extracted
21
22 using DCM and H₂O with 1% formic acid. Air-dried cyclic peptides were dissolved in MeOH and completely
23
24 dried under vacuum overnight.

25
26
27
28 For final cleavage, each Air-dried linear or cyclic peptides were dissolved in 3 mL of cleavage cocktail (95% v/v
29
30 TFA, 2.5% v/v triisopropylsilane and 2.5% v/v H₂O) for 1.5 hrs. Then, to precipitate the peptide, a cold mixture
31
32 of diethyl ether:hexane (1:1) was added and kept for 10 min at -20 °C. Peptide pellets were collected by
33
34 centrifuging (2,500 g) for 5 min, redissolved in 5 mL of methanol and dried under vacuum overnight.

35 36 37 **Peptide purification by HPLC**

38
39
40
41 Crude synCLP2-L, synCLP2-C1, synCLP2-C7 (chisopeptin) and chisopeptin congeners were purified on a Xbridge
42
43 Prep C18 HPLC column using a dual solvent system (A/B: H₂O/acetonitrile) (Figure S1). All synthetic peptides
44
45 were eluted using a linear gradient from 20 to 45% gradient of B. The identity of each linear or cyclic peptide
46
47 was confirmed by HRMS (Table S2 and Figure S2-4). Furthermore, ¹H, ¹³C, DEPT135, COSY, HSQC and HMBC
48
49 NMR spectra were recorded for chisopeptin and its five congeners (Figure S5-40). Structural assignments were
50
51 performed with additional information from NMR experiments (Table S3-8).

52 53 54 **Antimicrobial assays against Gram-positive bacteria, Gram-negative bacteria and fungi**

1
2
3
4
5 All antimicrobial assays were performed in 96-well microtiter plates using a broth micro-dilution method. For
6
7 fungi strains, overnight cultures were diluted 2,000-fold in YPD broth. Overnight cultures were diluted 1,000-
8
9 and 10,000-fold in LB broth for *Enterococcus faecium* and *Staphylococcus aureus*, respectively. For other
10
11 bacteria, overnight cultures were diluted 5,000-fold in LB broth. 100 μ L of each diluted culture was mixed with
12
13 100 μ L of LB broth containing a synthetic peptide at 2-fold serial dilutions across a 96-well microtiter plate row.
14
15 The final concentration of each synthetic peptide ranged from 64 to 0.125 μ g/mL. Plates were incubated at
16
17 37 $^{\circ}$ C (bacteria) or 30 $^{\circ}$ C (fungi) for 16 hrs. The lowest concentration that inhibited visible bacterial or fungal
18
19 growth was recorded as the minimum inhibition concentration (MIC). All MIC assays were performed in
20
21 duplicate (n = 2).
22
23
24

25 **Antibiotic resistant mutant selection**

26
27
28 A single *K. pneumoniae* ATCC 10031 colony was grown overnight at 37 $^{\circ}$ C in liquid LB medium. A portion of the
29
30 overnight culture was diluted to $\sim 10^8$ cells per microliter, which were plated into solid LB with chisopeptin at
31
32 4x its MIC at 5 μ L per well. The MICs of three individual appeared colonies were then determined using the
33
34 broth micro-dilution method described above. Genome was extracted from cultures of colonies that showed
35
36 an elevated MIC relative to the wild-type, and the resulting genome was sequenced by the MGI's DNBSEQ-T7
37
38 platform. Single-nucleotide polymorphisms (SNPs) for chisopeptin were identified using SNIPPY by mapping
39
40 DNBSEQ-seq reads to the reference genome of *K. pneumoniae* ATCC 10031.
41
42
43

44 **Cytotoxicity assessment**

45
46
47 The human HeLa cell line was grown at 37 $^{\circ}$ C in a 5% CO₂ atmosphere in DMEM. To allow HeLa cells to adhere
48
49 96-well flat bottom microtiter plates, they were incubated in DMEM at 37 $^{\circ}$ C for 24 hrs. Then, the DMEM
50
51 medium was replaced with 100 μ L of fresh DMEM medium containing each synthetic peptide at 10 serially
52
53 diluted concentrations ranging from 64 to 0.125 μ g/mL. After 48 hrs at 37 $^{\circ}$ C, 10 μ L of a CCK-8 solution was
54
55 added into each well. After 2 hrs at 37 $^{\circ}$ C, the absorbance at 450 nm for each well was detected for measuring
56
57
58

1
2
3
4 the cytotoxicity of each synthetic peptide using a microplate reader. Taxol and DMSO were used as the positive
5
6 and negative controls for the measurement of IC_{50} of each synthetic peptide against the human cell HeLa,
7
8 respectively.
9
10

11 **Results and Discussion**

12 **Prediction and synthesis of the CLP2 BGC products**

13
14
15
16
17
18 More than 90% sequenced bacterial BGCs are not or lowly expressed under standard laboratory conditions,
19
20 which could serve as rich sources for novel antibiotic discovery.²⁰⁻²³ With the increasing accuracy of NP
21
22 structure prediction algorithms and the advancement of modern synthetic organic chemistry, a bioactive
23
24 compound could be rapidly generated by structure prediction followed by chemical synthesis.^{24,25} The
25
26 synthetic-bioinformatic natural product (syn-BNP) approach has been widely used for *de novo* and targeted
27
28 discovery of diverse NRPs with novel structures or MOAs.²⁶⁻²⁹ In this study, we used this syn-BNP approach to
29
30 access linear or cyclic lipopeptide structures inspired by the CLP2 BGC for the test of antimicrobial activity. As
31
32 shown in Figure 2a, all genes in the CLP2 BGC was predicted to form a single operon considering their
33
34 expression were found to follow the same direction. The absence of any tailoring enzyme-encoding genes
35
36 within 10 kb of the CLP2 NRPS genes suggested that the product of the CLP2 BGC could not be post-modified
37
38 (Figure 2a and S39). Meanwhile, A-domain substrate specificity analysis allowed us to predict the building block
39
40 incorporated by each A-domain found in the CLP2 BGC with high accuracy (Figure 2a and Table S9). Therefore,
41
42 the linear peptide encoded by the CLP2 BGC was predicted to be the direct precursor to the final peptide.
43
44 Meanwhile, considering that the two amino acids, D -Dap-1 and D -Dap-7, in predicted peptides, could serve as
45
46 nucleophiles for cyclization of the predicted peptide through the C-terminal carboxyl, predicted CLP2 products
47
48 could appear as either linear or cyclic structures. Meanwhile, the condensation starter (Cs) domain that usually
49
50 couples a fatty acid onto the first amino acid of peptide were encoded by the CLP2 BGC (Figure S41). Among
51
52 the known bacterial CLPs, tetradecanoic acid is one of the most common lipids, so all predicted CLP2 peptides
53
54
55
56
57
58

(Table 1). These results indicated that synCLP2-C7 we named chisopeptin had relatively narrow antimicrobial spectrum. Naturally occurring bacterial lipopeptides appear as either linear or cyclic structures.^{6,7,31} It is very rare that one non-ribosomal lipopeptide BGC could encode both cyclic and linear peptides.^{6,7,31,32} Considering

Table 1 | Activities of synCLP2-L, synCLP2-C1, chisopeptin and its eight congeners against microorganisms and human cell (MIC: $\mu\text{g}/\text{mL}$). Colistin and amphotericin B were used as the positive controls. ND: not detected.

Type	Pathogens	synCLP2-L	synCLP2-C1	Chisopeptin	Chisopeptin-NF	Chisopeptin-HA	Chisopeptin-DA	Chisopeptin-V11	Chisopeptin-V9	Chisopeptin-L6-V9	Chisopeptin-L6-F9	Chisopeptin-A4-L6-F9	Colistin	Amphotericin B
Gram-negative bacteria	<i>Escherichia coli</i> K12	>64	>64	4	>64	>64	4	4	8	4	2	4	2	ND
	<i>Escherichia coli</i> DH5 α	>64	>64	8	>64	>64	8	4	8	8	4	8	<0.125	ND
	<i>Escherichia coli</i> ATCC 25922	>64	>64	4	>64	>64	8	8	8	4	2	8	0.5	ND
	<i>Klebsiella pneumoniae</i> ATCC 10031	>64	>64	4	>64	>64	4	4	8	8	2	4	0.5	ND
	<i>Klebsiella pneumoniae</i> RJA166	>64	>64	8	>64	>64	8	8	8	8	4	8	<0.125	ND
	<i>Klebsiella pneumoniae</i> RJF293	>64	>64	8	>64	>64	8	8	8	8	4	8	<0.125	ND
	<i>Klebsiella pneumoniae</i> RJF999	>64	>64	8	>64	>64	8	8	4	8	4	4	<0.125	ND
	<i>Erwinia amylovora</i> LXX	>64	>64	4	>64	64	4	4	4	4	2	2	0.25	ND
	<i>Vibrio cholerae</i> ATCC 39315	>64	32	4	>64	>64	8	8	4	8	2	4	>64	ND
	<i>Acinetobacter baumannii</i> ATCC 19606	>64	32	>64	>64	>64	64	>64	>64	>64	>64	32	1	ND
<i>Pseudomonas aeruginosa</i> ATCC 15692	>64	>64	32	>64	>64	16	>64	>64	>64	>64	32	>64	0.5	ND
Gram-positive bacteria	<i>Bacillus subtilis</i> ATCC 23857	32	4	4	>64	16	4	2	2	4	2	4	>64	ND
	<i>Enterococcus faecalis</i> ATCC 29212	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	ND
	<i>Staphylococcus aureus</i> USA300	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	ND
	<i>Mycobacterium smegmatis</i> mc2 155	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	ND
Fungi	<i>Candida albicans</i> ATCC 10231	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	0.5
	<i>Saccharomyces cerevisiae</i> BY4741	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	>64	0.25
Human cell line	HeLa	ND	ND	>64	ND	ND	ND	>64	>64	>64	>64	>64	>64	ND

only chisopeptin showed good anti-Gram-negative activity and either synCLP2-L or synCLP2-C1 was not active against tested pathogens, we speculate that chisopeptin is likely to be closest to natural product that the CLP2 BGC encodes (Figure 2d). The structure of chisopeptin was further confirmed using HRMS and 1D/2D NMR data (Table S2, Figure S1-2 and 5-10). Then, we synthesized three chisopeptin congeners by changing the length of fatty acid side chain, including chisopeptin-NF without fatty acid side chain, chisopeptin-HA with the hexanoic acid as the fatty acid side chain and chisopeptin-DA with the decanoic acid as the fatty acid side chain (Figure S1 and S3). As shown in Table 1, neither chisopeptin-NF nor chisopeptin-HA was active against tested Gram-negative bacteria and chisopeptin-DA showed compared anti-Gram-negative activity to chisopeptin

(tetradecanoic acid as the fatty acid side chain). The results indicate that a long fatty acid side chain is necessary to the activity of chisopeptin against Gram-negative bacteria.

Prediction and synthesis of the five chisopeptin congener BGC products




Figure 3 | Discovery of five chisopeptin congeners. a, The BGCs of CLP2 and five chisopeptin congeners. b, Linear peptide sequences of chisopeptin and its five congeners. c, MIC data against Gram-negative or Gram-positive pathogens for chisopeptin and its five congeners (n=2).

In our first-round search pipeline to discover novel CLP BGCs using at least two PCAAs in a peptide as the motif, only one representative CLP BGC among highly similar BGCs was chosen for next-step study. Therefore, we re-searched the predicted NRP database from 42 underexplored Gram-negative bacterial genera and found the five potential chisopeptin congener BGCs. Intriguingly, all the five BGCs are from the genus *Pedobacter*, which

1
2
3
4 has not traditionally been part of microbial NP discovery pipelines.¹³ Similar to the CLP2 BGC, the expression
5 of all genes in each BGC were also found to follow the same direction, which possibly form a single operon
6 (Figure 3a and S42-46). Except for a MBL fold metallo-hydrolase encoded by the chisozeptin-L6-V9 BGC, there
7 aren't any tailoring enzymes encoded by the five BGCs (Figure 3a and S41-45). Through A-domain substrate
8 specificity analysis, we found that there were one to three different AAs in linear peptides predicted by the five
9 BGCs from chisozeptin (Table S10-14, Figure 3b and S1). Then, all five chisozeptin congeners, including
10 chisozeptin-V11, chisozeptin-V9, chisozeptin-L6-V9, chisozeptin-L6-F9 and chisozeptin-A4-L6-F9, were
11 designed to be cyclized through the hydroxyl of _D-Dap-7 and rapidly generated by solid-phase peptide synthesis.
12 Their structures were further confirmed using HRMS and 1D/2D NMR data (Table S2 and S4-8, Figure S4 and
13 S11-40). Similar to chisozeptin, all five congeners showed similar antimicrobial spectra to chisozeptin and were
14 active against four Gram-negative pathogens with MIC from 2 to 8 μg/mL. In a subset of these congeners,
15 antibacterial activities of chisozeptin-L6-F9 in particular, were two-fold higher than those of chisozeptin
16 (Figure 3c and Table 1). Collectively, a novel family of cationic cyclolipopeptides, including chisozeptin and its
17 five congeners, were identified with good activities against a panel of Gram-negative pathogens.

34 35 **Mode of action studies**

36
37
38 Then, we tested whether *K. pneumoniae* ATCC 10031 would be lysed using 4× the MIC of chisozeptin, and
39 found that chisozeptin could not result in rapid bacteriolysis (Figure 4a). Considering that chisozeptin as a
40 cationic cyclolipopeptide might target LPS in the outer membrane of Gram-negative pathogens, we assayed
41 for the ability of LPS to suppress chisozeptin to explore MOA of chisozeptin. However, when LPS was added to
42 the LB medium, the MIC of chisozeptin did not increase in a dose-dependent manner for *K. pneumoniae*, which
43 indicated that chisozeptin could not target LPS and its anti-Gram-negative activity was not dependent on LPS
44 (Figure 4b). Then, we tried to raise mutants to chisozeptin and selected *K. pneumoniae* mutants that could
45 grow on 4× MIC of chisozeptin. At 4× chisozeptin's MIC resistant mutants appeared at a frequency of 1×10⁻⁶.
46 We sequenced the genomes of three representative resistant strains (Figure 4c). In all cases, the resistant
47
48
49
50
51
52
53
54
55
56
57
58

strains contained a mutation (in-frame or frame-shift deletion) in the gene *yejM* and no other mutations were detected in the genome (Figure 4d). When the gene *yejM* under the expression of the constitutive promoter *rpsLp* was complemented to the three *K. pneumoniae* mutants, all of them became sensitive to chisopeptin again (Figure




Figure 4 | Mode-of-action of chisopeptin against the strain *K. pneumoniae* ATCC 10031. a, Bactericidal activity of chisopeptin against *K. pneumoniae*. Cultures were incubated with each antibiotic at 4xits MIC. b, The *K. pneumoniae* antibacterial activity of chisopeptin was determined in the presence of different concentrations of LPS (n = 2). c, The MICs (µg/mL) of chisopeptin against *K. pneumoniae* mutants (n = 2). Colistin was used as the positive control. d, The mutated results of the gene *yejM* in the three chisopeptin resistant *K. pneumoniae* strains. e, The MICs (µg/mL) of chisopeptin against *K. pneumoniae* mutants with pET28a or pET-*yejM* (n = 2). Colistin was used as the positive control. f, A proposed model-of-action of chisopeptin against Gram-negative pathogens (i.e., *K. pneumoniae*).

4e). YejM, as an inner membrane protein, is a key regulator of outer membrane integrity and LPS biogenesis by directly detecting LPS on the periplasmic leaflet of the inner membrane in Gram-negative bacteria.^{33,34} In

1
2
3
4 detail, YejM as an anti-adaptor protein can compete with FtsH for LapB to inhibit LpxC degradation (Figure
5 4f).³⁵ Deletion of the gene *yejM* could promote LpxC degradation, thus resulting in reduced LPS levels.^{34,36}
6
7 Previous studies indicated that the mutation of *yejM* also led to higher resistance to chloramphenicol in *E.*
8
9 *coli*.³⁷ Considering that chisopeptin did not directly target LPS, we proposed that chisopeptin might target the
10
11 inner membrane protein YejM, thus interrupting normal LPS biogenesis and exhibiting anti-Gram-negative
12
13 activity (Figure 4f). To explore whether chisopeptin could directly bind to YejM *in vitro*, we tried to express and
14
15 purify YejM but failed, which might be because YejM is a membrane protein (data not shown). The more
16
17 detailed MOA for chisopeptin could be further explored in the future.
18
19
20
21
22

23 Conclusion

24
25 Infections caused by multidrug-resistant Gram-negative pathogens are threatening to overwhelm
26
27 healthcare systems worldwide.^{2,3} In recent years, underexplored Gram-negative bacteria exhibit large
28
29 biosynthetic potential for the discovery of novel antimicrobial peptides.¹²⁻¹⁵ In this study, we used the two-
30
31 round motif search pipelines, including at least two PCAAs and multiple D-AAs, to screen >8900 underexplored
32
33 Gram-negative bacterial genomes and identified a novel family of cationic cyclolipoptides (chisopeptin and
34
35 its five congeners) with good activities against a panel of Gram-negative pathogens. All six synBNPs have very
36
37 unique structural scaffolds, which contain six PCAAs and nine D-AAs out of 12 building blocks. Noteworthy, the
38
39 kind of cationic cyclolipoptides was identified to have a potential MOA to kill *K. pneumoniae* by possibly
40
41 targeting the inner membrane protein YejM that regulates LPS biogenesis. Considering that *K. pneumoniae* has
42
43 relatively high frequency to generate resistant mutants to chisopeptin, combinations of chisopeptin-type
44
45 antibiotics and of known Gram-negative-active drugs with different MOAs (i.e., penicillin-binding protein-
46
47 targeting meropenem, ribosome-targeting amikacin and DNA-targeting moxifloxacin), provide an alternative
48
49 strategy for further pre-clinical development of this novel family of cationic cyclolipoptides.³⁸ As seen with
50
51 the characterization of bacterial CLPs, the study of chisopeptin-type structures inspired by underexplored BGC
52
53 families provides an alternative approach to develop novel CLP-based therapeutics with diverse MOAs.^{39,40}
54
55
56
57
58
59
60

Supporting Information

Supporting Information is available and includes chisopeptin and its congeners domain prediction tables, chisopeptin and its congeners BGC analysis, chisopeptin and its congeners HRMS and NMR spectra.

Conflict of Interest

There is no conflict of interest to report.

Funding Information

This work was supported by the National Key Research and Development Program of China (2023YFA0916200 and 2023YFA0914200), the National Natural Science Foundation of China (32370070) and Shanghai Municipal Science and Technology Major Project.

Acknowledgments

We thank Prof. Jianming Zhang and Xiaowen Wang at Shanghai Jiao Tong University for cytotoxicity assessment.

References

1. Abbas, A.; Barkhouse, A.; Hackenberger, D.; Wright, G. D. Antibiotic resistance: A key microbial survival mechanism that threatens public health. *Cell Host Microbe* **2024**, *32*, 837-851.
2. GBD 2019 Antimicrobial Resistance Collaborators. Global mortality associated with 33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* **2022**, *400*, 2221-2248.
3. Zowawi, H. M.; Harris, P. N.; Roberts, M. J.; Tambyah, P. A.; Schembri, M. A.; Pezzani, M. D.; Williamson, D. A.; Paterson, D. L. The emerging threat of multidrug-resistant Gram-negative bacteria in urology. *Nat. Rev. Urol.* **2015**, *12*, 570-584.
4. Lewis, K.; Lee, R. E.; Brötz-Oesterhelt, H.; Hiller, S.; Rodnina, M. V.; Schneider, T.; Weingarh, M.; Wohlgemuth, I. Sophisticated natural products as antibiotics. *Nature* **2024**, *632*, 39-49.

- 1
2
3
4
5 5. Atanasov, A. G.; Zotchev, S. B.; Dirsch, V. M.; International Natural Product Sciences Taskforce; Supuran, C.
6
7 T. Natural products in drug discovery: advances and opportunities. *Nat. Rev. Drug Discov.* **2021**, *20*, 200-216.
8
9
10 6. Liu, Y.; Ding, S.; Shen, J.; Zhu, K. Nonribosomal antibacterial peptides that target multidrug-resistant bacteria.
11
12 *Nat. Prod. Rep.* **2019**, *36*, 573-592.
13
14
15 7. Chen, J.; Wang, W.; Hu, X.; Yue, Y.; Lu, X.; Wang, C.; Wei, B.; Zhang, H.; Wang, H. Medium-sized peptides
16
17 from microbial sources with potential for antibacterial drug development. *Nat. Prod. Rep.* **2024**, *41*, 1235-1263.
18
19
20 8. Fetse, J.; Kandel, S.; Mamani, U. F.; Cheng, K. Recent advances in the development of therapeutic peptides.
21
22 *Trends Pharmacol. Sci.* **2023**, *44*, 425-441.
23
24
25 9. Biswas, S.; Brunel, J. M.; Dubus, J. C.; Reynaud-Gaubert, M.; Rolain, J. M. Colistin: an update on the antibiotic
26
27 of the 21st century. *Expert Rev. Anti-Infect.* **2012**, *10*, 917-934.
28
29
30 10. Li, Y. X.; Zhong, Z.; Zhang, W. P.; Qian, P. Y. Discovery of cationic nonribosomal peptides as Gram-negative
31
32 antibiotics through global genome mining. *Nat. Commun.* **2018**, *9*, 3273.
33
34
35 11. Wang, Z.; Koirala, B.; Hernandez, Y.; Zimmerman, M.; Park, S.; Perlin, D. S.; Brady, S. F. A naturally inspired
36
37 antibiotic to target multidrug-resistant pathogens. *Nature* **2022**, *601*, 606-611.
38
39
40 12. Li, L. Accessing hidden microbial biosynthetic potential from underexplored sources for novel drug
41
42 discovery. *Biotechnol. Adv.* **2023**, *66*, 108176.
43
44
45 13. Masschelein, J.; Jenner, M.; Challis, G. L. Antibiotics from Gram-negative bacteria: a comprehensive
46
47 overview and selected biosynthetic highlights. *Nat. Prod. Rep.* **2017**, *34*, 712-783.
48
49
50 14. Brinkmann, S.; Spohn, M. S.; Schäberle, T. F. Bioactive natural products from Bacteroidetes. *Nat. Prod. Rep.*
51
52 **2022**, *39*, 1045-1065.
53
54
55
56
57
58

- 1
2
3
4
5 15. Brinkmann, S.; Kurz, M.; Patras, M. A.; Hartwig, C.; Marner, M.; Leis, B.; Billion, A.; Kleiner, Y.; Bauer, A.;
6
7 Toti, L.; Pöverlein, C.; Hammann, P. E.; Vilcinskas, A.; Glaeser, J.; Spohn, M.; Schäberle, T. F. Genomic and
8
9 chemical decryption of the bacteroidetes phylum for its potential to biosynthesize natural products. *Microbiol.*
10
11 *Spectr.* **2022**, *10*, e0247921.
12
13
14 16. Sun, R.; You, R.; Yu, X.; Zhao, D.; Li, L. Discovery and synthesis of a Gram-negative-active cationic lipopeptide
15
16 antibiotic inspired by primary sequences from underexplored Gram-negative bacteria. *Org. Lett.* **2024**, *26*,
17
18 1348-1352.
19
20
21 17. Zhang, H.; Chen, S. Cyclic peptide drugs approved in the last two decades (2001-2021). *RSC Chem. Biol.*
22
23 **2021**, *3*, 18-31.
24
25
26 18. Armstrong, D. W.; Berthod, A. Occurrence of D-amino acids in natural products. *Nat. Prod. Bioprospect.*
27
28 **2023**, *13*, 47.
29
30
31 19. Abdulbagi, M.; Wang, L.; Siddig, O.; Di, B.; Li, B. D-amino acids and D-amino acid-containing peptides:
32
33 potential disease biomarkers and therapeutic targets? *Biomolecules* **2021**, *11*, 1716.
34
35
36 20. Rutledge, P. J.; Challis, G. L. Discovery of microbial natural products by activation of silent biosynthetic gene
37
38 clusters. *Nat. Rev. Microbiol.* **2015**, *13*, 509-523.
39
40
41 21. Gavriilidou, A.; Kautsar, S. A.; Zaburannyi, N.; Krug, D.; Müller, R.; Medema, M. H.; Ziemert, N. Compendium
42
43 of specialized metabolite biosynthetic diversity encoded in bacterial genomes. *Nat. Microbiol.* **2022**, *7*, 726-
44
45 735.
46
47
48 22. Medema, M. H.; de Rond, T.; Moore, B. S. Mining genomes to illuminate the specialized chemistry of life.
49
50
51 *Nat. Rev. Genet.* **2021**, *22*, 553-571.
52
53
54
55
56
57
58

- 1
2
3
4
5 23. Hemmerling, F.; Piel, J. Strategies to access biosynthetic novelty in bacterial genomes for drug discovery.
6
7 *Nat. Rev. Drug Discov.* **2022**, *21*, 359-378.
8
9
10 24. Blin, K.; Shaw, S.; Augustijn, H. E.; Reitz, Z. L.; Biermann, F.; Alanjary, M.; Fetter, A.; Terlouw, B. R.; Metcalf,
11
12 W. W.; Helfrich, E. J. N.; van Wezel, G. P.; Medema, M. H.; Weber, T. antiSMASH 7.0: new and improved
13
14 predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids Res.* **2023**, *51*, W46-
15
16 W50.
17
18
19 25. Stachelhaus, T.; Mootz, H. D.; Marahiel, M. A. The specificity-conferring code of adenylation domains in
20
21 nonribosomal peptide synthetases. *Chem. Biol.* **1999**, *6*, 493-505.
22
23
24
25 26. Chu, J.; Vila-Farres, X.; Inoyama, D.; Ternei, M.; Cohen, L. J.; Gordon, E. A.; Reddy, B. V.; Charlop-Powers, Z.;
26
27 Zebroski, H. A.; Gallardo-Macias, R.; Jaskowski, M.; Satish, S.; Park, S.; Perlin, D. S.; Freundlich, J. S.; Brady, S. F.
28
29 Discovery of MRSA active antibiotics using primary sequence from the human microbiome. *Nat. Chem. Biol.*
30
31 **2016**, *12*, 1004-1006.
32
33
34 27. Chu, J.; Vila-Farres, X.; Brady, S. F. Bioactive synthetic-bioinformatic natural product cyclic peptides inspired
35
36 by nonribosomal peptide synthetase gene clusters from the human microbiome. *J. Am. Chem. Soc.* **2019**, *141*,
37
38 15737-15741.
39
40
41
42 28. Wang, Z.; Koirala, B.; Hernandez, Y.; Zimmerman, M.; Brady, S. F. Bioinformatic prospecting and synthesis
43
44 of a bifunctional lipopeptide antibiotic that evades resistance. *Science* **2022**, *376*, 991-996.
45
46
47 29. Li, L.; Koirala, B.; Hernandez, Y.; MacIntyre, L. W.; Ternei, M. A.; Russo, R.; Brady, S. F. Identification of
48
49 structurally diverse menaquinone-binding antibiotics with *in vivo* activity against multidrug-resistant
50
51 pathogens. *Nat. Microbiol.* **2022**, *7*, 120-131.
52
53
54
55
56
57
58

- 1
2
3
4
5 30. Rausch, C.; Hoof, I.; Weber, T.; Wohlleben, W.; Huson, D. H. Phylogenetic analysis of condensation domains
6
7 in NRPS sheds light on their functional evolution. *BMC Evol. Biol.* **2007**, *7*, 78.
8
9
10 31. Zhang, S.; Chen, Y.; Zhu, J.; Lu, Q.; Cryle, M. J.; Zhang, Y.; Yan, F. Structural diversity, biosynthesis, and
11
12 biological functions of lipopeptides from *Streptomyces*. *Nat. Prod. Rep.* **2023**, *40*, 557-594.
13
14
15 32. Götze, S.; Vij, R.; Burow, K.; Thome, N.; Urvat, L.; Schlosser, N.; Pflanze, S.; Müller, R.; Hänsch, V. G.;
16
17 Schlabach, K.; Fazlikhani, L.; Walther, G.; Dahse, H. M.; Regestein, L.; Brunke, S.; Hube, B.; Hertweck, C.; Franken,
18
19 P.; Stallforth, P. Ecological niche-inspired genome mining leads to the discovery of crop-protecting
20
21 nonribosomal lipopeptides featuring a transient amino acid building block. *J. Am. Chem. Soc.* **2023**, *145*, 2342-
22
23 2353.
24
25
26 33. Guest, R. L.; Rutherford, S. T.; Silhavy, T. J. Border control: regulating LPS biogenesis. *Trends Microbiol.* **2021**,
27
28 *29*, 334-345.
29
30
31
32 34. Clairfeuille, T.; Buchholz, K. R.; Li, Q.; Verschueren, E.; Liu, P.; Sangaraju, D.; Park, S.; Noland, C. L.; Storek,
33
34 K. M.; Nickerson, N. N.; Martin, L.; Dela Vega, T.; Miu, A.; Reeder, J.; Ruiz-Gonzalez, M.; Swem, D.; Han, G.;
35
36 DePonte, D. P.; Hunter, M. S.; Gati, C.; Shahidi-Latham, S.; Xu, M.; Skelton, N.; Sellers, B. D.; Skippington, E.;
37
38 Sandoval, W.; Hanan, E. J.; Payandeh, J.; Rutherford, S. T. Structure of the essential inner membrane
39
40 lipopolysaccharide-PbgA complex. *Nature* **2020**, *584*, 479-483.
41
42
43 35. Shu, S.; Mi, W. Regulatory mechanisms of lipopolysaccharide synthesis in *Escherichia coli*. *Nat. Commun.*
44
45 **2022**, *13*, 4576.
46
47
48 36. Fivenson, E. M.; Bernhardt, T. G. An essential membrane protein modulates the proteolysis of LpxC to
49
50 control lipopolysaccharide synthesis in *Escherichia coli*. *mBio* **2020**, *11*, e00939-20.
51
52
53
54
55
56
57
58

- 1
2
3
4
5 37. Duo M, Hou S, Ren D. Identifying *Escherichia coli* genes involved in intrinsic multidrug resistance. *Appl.*
6
7 *Microbiol. Biotechnol.* **2008**, *81*, 731-41.
8
9
10 38. Tyers, M., Wright, G. S. Drug combinations: a strategy to extend the life of antibiotics in the 21st century.
11
12 *Nat. Rev. Microbiol.* **2019**, *17*, 141-155.
13
14
15 39. Zhao, S.; Adamiak, J. W.; Bonifay, V.; Mehla, J.; Zgurskaya, H. I.; Tan, D. S. Defining new chemical space for
16
17 drug penetration into Gram-negative bacteria. *Nat. Chem. Biol.* **2020**, *16*, 1293-1302.
18
19
20 40. Miethke, M.; Pieroni, M.; Weber, T.; Brönstrup, M.; Hammann, P.; Halby, L.; Arimondo, P. B.; Glaser, P.;
21
22 Aigle, B.; Bode, H. B.; Moreira, R.; Li, Y.; Luzhetskyy, A.; Medema, M. H.; Pernodet, J. L.; Stadler, M.; Tormo, J.
23
24 R.; Genilloud, O.; Truman, A. W.; Weissman, K. J.; Takano, E.; Sabatini, S.; Stegmann, E.; Brötz-Oesterhelt, H.;
25
26 Wohlleben, W.; Seemann, M.; Empting, M.; Hirsch, A. K. H.; Loretz, B.; Lehr, C. M.; Titz, A.; Herrmann, J.; Jaeger,
27
28 T.; Alt, S.; Hesterkamp, T.; Winterhalter, M.; Schiefer, A.; Pfarr, K.; Hoerauf, A.; Graz, H.; Graz, M.; Lindvall, M.;
29
30 Ramurthy, S.; Karlén, A.; van Dongen, M.; Petkovic, H.; Keller, A.; Peyrane, F.; Donadio, S.; Fraise, L.; Piddock,
31
32 L. J. V.; Gilbert, I. H.; Moser, H. E.; Müller, R. Towards the sustainable discovery and development of new
33
34 antibiotics. *Nat. Rev. Chem.* **2021**, *5*, 726-749
35
36
37

38 **Table of Contents Graphic**

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

